

DEFORMATIONS OF LINEAR POISSON ORBIFOLDS

GILLES HALBOUT, JEAN-MICHEL OUDOM, AND XIANG TANG

ABSTRACT. Let Γ be a finite group acting faithfully and linearly on a vector space V . Let $T(V)$ ($S(V)$) be the tensor (symmetric) algebra associated to V which has a natural Γ action. We study generalized quadratic relations on the tensor algebra $T(V) \rtimes \Gamma$. We prove that the quotient algebras of $T(V) \rtimes \Gamma$ by such relations satisfy PBW property. Such quotient algebras can be viewed as quantizations of linear or constant Poisson structures on $S(V) \rtimes \Gamma$, and are natural generalizations of symplectic reflection algebras.

1. INTRODUCTION

Poisson structure on a manifold is a bivector field π whose Schouten-Nijenhuis bracket with itself vanishes, i.e. $\pi \in \Gamma(\wedge^2 TM)$, and $[\pi, \pi] = 0$. The problem of deformation quantization of a Poisson manifold was solved by Kontsevich in his semina paper [10]. In this paper, we study quantization problem of Poisson structures on an orbifold following [9].

The study of the first and third author in [9] starts with the idea that Poisson structures on an algebra A should correspond to infinitesimal deformations of A . According to Gerstenhaber's theory, an infinitesimal deformation of an algebra is classified by a second Hochschild cohomology class in $H^2(A, A)$ whose Gerstenhaber bracket with itself is zero. This type of cohomology class is called a Poisson structure ([2], [15]) on A . Applying this idea to orbifold, we can represent an orbifold X by a proper étale groupoid $\mathcal{G}$ [11] (different representations of same orbifold are Morita equivalent as Lie groupoids). We consider the smooth groupoid algebra $C_c^\infty(\mathcal{G})$ associated to $\mathcal{G}$. We studied in [9] Poisson structures on $C_c^\infty(\mathcal{G})$. We find that Poisson structures on $C_c^\infty(\mathcal{G})$ are richer than we naturally expect from geometry. On an orbifold, multivector fields and Schouten-Nijenhuis bracket are well defined. Accordingly, we can consider bivector fields on X whose Schouten-Nijenhuis bracket with themselves vanish. We found that [9][Theorem 4.1] there are many more Poisson structures on $C_c^\infty(\mathcal{G})$ than the above type of bivector fields on X . For example, in the case of a finite group Γ acting on a symplectic vector space V , we [9][Corollary 4.2] find Poisson structures on $S(V^*) \rtimes \Gamma$ which have supports on codimension 2 fixed point subspaces, where $S(V^*)$ is the algebra of real coefficients symmetric polynomials on the dual vector space V^* .

In this paper, we continue our study of Poisson structures in the above framework. We will study Poisson structures in a neighborhood of a point in a reduced orbifold. Locally, a reduced orbifold can always be viewed as a quotient of a finite group acting faithfully and linearly on an open set of $\mathbb{R}^n$. This leads us to study the following data. Let Γ be a finite group acting on a vector space V faithfully, and $S(V^*)$ be the algebra of polynomials on V^* . The Γ action on V^* defines the crossed product algebra $S(V^*) \rtimes \Gamma$. According to [13], the second Hochschild cohomology $H^2(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma)$ is equal to

$$(S(V^*) \otimes \wedge^2 V)^\Gamma \oplus \left(\sum_{\gamma \in \Gamma, l(\gamma)=2} S(V^{\gamma^*}) \otimes \wedge^2 N^\gamma \right)^\Gamma.$$

In the above equation, $l(\gamma)$ is the codimension of the γ fixed point subspace V^γ and N^γ consists of vectors in V vanishing on V^{γ^*} (the fixed points subspace of γ action on V^*), and Γ acts on the set $S := \{\gamma \in \Gamma, l(\gamma) = 2\}$ by conjugation.

We study two types of Poisson structures on $S(V^*) \rtimes \Gamma$ which are of the forms

$$i) \text{ Hom}(\wedge^2 V^*, \mathbb{R}\Gamma), \quad ii) \text{ Hom}(\wedge^2 V^*, V^* \otimes_{\mathbb{R}} \mathbb{R}\Gamma).$$

The first type of Poisson structure can be viewed as constant value Poisson structures, and the second type can be viewed as linear Poisson structures, which define generalized ‘‘Lie’’ algebra structures on

$V^* \otimes_{\mathbb{R}} \mathbb{R}\Gamma$. Our main theorems for these Poisson structures are that the quotient algebras of $T(V^*) \rtimes \Gamma$ by the relations defined by the above two types of Poisson structures satisfy PBW property. (For general linear Poisson structures, we need to assume that V is equipped with a Γ -invariant complex structure.) The way we prove such theorems is using the Braverman-Gaitsory conditions [3] for PBW property. However, the proof for the second type is quite involved. We need to use properties of finite subgroups of $GL(2, \mathbb{C})$, which is closely related to McKay correspondence. A new and interesting phenomena we found in the proof of PBW property is that we have to have a nontrivial coboundary term for the bracket $[\pi, \pi]$ of the linear Poisson structure π . This kind of term never shows up in the study of PBW property for Lie algebras and symplectic reflection algebras. The PBW property of the quotient algebras shows that they define quantizations of the Poisson structures on $S(V^*) \rtimes \Gamma$. This confirms that any constant or linear Poisson structures on $S(V^*) \rtimes \Gamma$ can be quantized, and gives a strong evidence that the deformation theory of the algebra $S(V^*) \rtimes \Gamma$ is formal.

The second part of this paper is dedicated to studying various properties and examples of the above two types of Poisson structures and their quantizations. We mention a few of them here. Firstly, using Poisson cohomology computation, we are able to give a new computation of Hochschild cohomology of a symplectic reflection algebra [6][Theorem 1.8]. The advantage of our work is that our result works for Laurent series of $\hbar$ so that we can drop the assumption “except possibly a countable set” in [6][Theorem 1.8]. Secondly, assuming a finite group Γ acts faithfully and linearly on a Lie algebra $\mathfrak{g}$, we compute the Hochschild cohomology of $H^\bullet(\mathcal{U}(\mathfrak{g}) \rtimes \Gamma, \mathcal{U}(\mathfrak{g}) \rtimes \Gamma)$ with $\mathcal{U}(\mathfrak{g})$ the universal enveloping algebra of $\mathfrak{g}$. This is a natural generalization of results [1] for a finite group action on a symplectic vector space. Our result shows that the Hochschild cohomology $H^\bullet(\mathcal{U}(\mathfrak{g}) \rtimes \Gamma, \mathcal{U}(\mathfrak{g}) \rtimes \Gamma)$ is computed by the noncommutative Poisson cohomology associated to the Γ action and Lie-Poisson structure π on $\mathfrak{g}$. Thirdly, if V is equipped with a Γ invariant Lie Poisson structure, we introduce a large class of examples of linear Poisson structures on $S(V^*) \rtimes \Gamma$. Quantizations of these linear Poisson structures should be viewed as natural generalizations of symplectic reflection algebras. Analogous to the symplectic case, we are able to prove that the Poisson cohomology of such a general linear Poisson structure is determined by the data supported at the identity of Γ in the case that Γ is abelian.

In the third part of this paper, we restrict ourselves to $\mathbb{R}^2$ with a cyclic group $\mathbb{Z}_n$ action. Let ω be the standard symplectic form on $\mathbb{R}^2$ and π be the corresponding Poisson structure. We study the quantization of the Poisson structure $\pi_\gamma : x \wedge y \rightarrow \pi(x, y)\gamma$ on $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$ for $\gamma \in \mathbb{Z}_n$ (We identify $\mathbb{R}^{2*}$ with $\mathbb{R}^2$). Nadaud [12] gave a Moyal type formula for such a deformation quantization. Here, we use this formula to study the center of the quantization. Our computation shows that the center of the quantization of π_γ is not isomorphic to the center of the algebra $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$. Instead, the center of the quantization is a nontrivial deformation of the center of $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$. This suggests that there is no analog of Duflo’s isomorphism for the quantization of the Poisson structure π_γ , and also that deformation quantization of π_γ on $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$ is closely connected to the deformation of the underlying orbifold singularity. We plan to study the relation between deformations of $S(V^*) \rtimes \Gamma$ and deformations of the underlying orbifold V/Γ in the near future.

This paper is organized as follows. In the second section, we review some results about Hochschild cohomology $H^\bullet(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma)$ in [13] and [9], and also the Braverman-Gaitsory conditions for PBW property [3]; in the third section, we prove that constant and linear Poisson structures on $S(V^*) \rtimes \Gamma$ can be quantized; in the forth section, we study various properties and examples of constant and linear Poisson structures on $S(V^*) \rtimes \Gamma$; in the fifth section, using Nadaud’s formula, we study the centers of quantizations of some Poisson structures on $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$.

Acknowledgements: We would like to thank Cédric Bonnafe for discussion about finite subgroups of $GL(2, \mathbb{C})$, Georges Pinczon for showing the results of Nadaud [12], and Victor Ginzburg for general discussion of symplectic reflection algebras and Duflo’s isomorphism. The research of the third author is partially supported by NSF Grant 0703775.

2. PRELIMINARIES AND NOTATIONS

In this whole paper, Γ is a finite group, acting faithfully and linearly on a finite dimensional real vector space V . We fix on V a Γ -invariant metric. We denote $C(\Gamma)$ the set of conjugacy classes of Γ . For any element γ in Γ , let V^γ be the γ -invariant subspace of V , N^γ be the subspace of V orthogonal to V^γ which is the direct sum of all nontrivial representations of $G(\gamma)$ (the subgroup of Γ generated by γ), $l(\gamma)$ be the real codimension of V^γ , and $Z(\gamma)$ the centralizer of γ in Γ . In this paper, we always work with the field $\mathbb{R}$. All dimensions, algebras, and tensor products if not specified are over the field $\mathbb{R}$. For the convenience of proofs, we are many times using the following complexification trick,

$$(1) \quad S_{\mathbb{R}}(V^*) \rtimes_{\mathbb{R}} \mathbb{C}\Gamma \cong S_{\mathbb{C}}(V^* \otimes \mathbb{C}) \rtimes_{\mathbb{C}} \mathbb{C}\Gamma \cong \left(S_{\mathbb{R}}(V^*) \rtimes_{\mathbb{R}} \mathbb{R}\Gamma \right) \otimes_{\mathbb{R}} \mathbb{C}.$$

This helps us to deduce the results in $\mathbb{R}$ from their complex versions where γ action is diagonalizable for any $\gamma \in \Gamma$. Many results in this paper hold true for field $\mathbb{C}$ and even more general field with characteristic 0 in which the order of Γ is invertible.

2.1. The Koszul complex and the Hochschild cohomology of $S(V^*) \rtimes \Gamma$. The algebra $S(V^*) \rtimes \Gamma$ is generated by V^* and Γ with the quadratic relations :

$$x \otimes y \otimes \gamma - y \otimes x \otimes \gamma, \quad \gamma \otimes x - {}^\gamma x \otimes \gamma,$$

for all x and y in V^* and γ in Γ , and ${}^\gamma x$ is the image of x under the γ action. Moreover, $S(V^*) \rtimes \Gamma$ is a Koszul algebra over the semi-simple algebra $\mathbb{R}\Gamma$. The general theory of Koszul algebras over a semi-simple algebra gives therefore a small complex which calculates the Hochschild cohomology of $S(V^*) \rtimes \Gamma$:

$$CK^\bullet(S(V^*) \rtimes \Gamma) = \left(\bigoplus_{\gamma \in \Gamma} S(V^*) \otimes \wedge^\bullet V \right)^\Gamma.$$

A n -cochain f of this complex splits in a sum of maps f_γ in $S(V^*) \otimes \wedge^n V$. The Γ -invariance can be written :

$$(2) \quad {}^g f_\gamma({}^{g^{-1}}x_1, \dots, {}^{g^{-1}}x_n) = f_{g\gamma g^{-1}}(x_1, \dots, x_n),$$

which explains that $CK^\bullet(S(V^*) \rtimes \Gamma)$ splits in a sum of sub-complexes :

$$CK^\bullet(S(V^*) \rtimes \Gamma) = \bigoplus_{\gamma \in C(\Gamma)} (S(V^*) \otimes \wedge^\bullet V)^{Z(\gamma)}$$

with the boundary

$$\partial_\gamma(f)(x_0, \dots, x_n) = \sum_{i=0}^n (-1)^i f(x_0, \dots, \widehat{x}_i, \dots, x_n)(x_i - {}^\gamma x_i),$$

for $x_0, \dots, x_n \in V^*$.

Using this small complex, Neumaier, Pflaum, Posthuma and the third author calculated in [13] the Hochschild cohomology of $S(V^*) \rtimes \Gamma$:

$$(3) \quad H^\bullet(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma) = \bigoplus_{\gamma \in C(\Gamma)} \left(S(V^{\gamma^*}) \otimes \wedge^{\bullet-l(\gamma)} V^\gamma \otimes \wedge^{l(\gamma)} N^\gamma \right)^{Z(\gamma)}.$$

This statement in $\mathbb{R}$ is easily deduced from its complex version [13] using the trick (1).

We point out that the projection $\text{pr}_\gamma : (S(V^*) \otimes \wedge^\bullet V)^{Z(\gamma)} \rightarrow (S(V^{\gamma^*}) \otimes \wedge^{\bullet-l(\gamma)} V^\gamma \otimes \wedge^{l(\gamma)} N^\gamma)^{Z(\gamma)}$ and the embedding $\iota : (S(V^{\gamma^*}) \otimes \wedge^{\bullet-l(\gamma)} V^\gamma \otimes \wedge^{l(\gamma)} N^\gamma)^{Z(\gamma)} \rightarrow (S(V^*) \otimes \wedge^\bullet V)^{Z(\gamma)}$ are inverse quasi-isomorphisms of complexes. Another useful remark is that if $l(\gamma)$ =dimension of N^γ is odd, the determinant of γ action on N^γ is -1 (otherwise γ has an eigenvalue 1 as γ is an isometry). Therefore $S(V^{\gamma^*}) \otimes \wedge^{\bullet-l(\gamma)} V^\gamma \otimes \wedge^{l(\gamma)} N^\gamma$ has no γ invariant element if $l(\gamma)$ is odd. Therefore, Poisson brackets on $S(V^*) \rtimes \Gamma$ do not contain γ -component for $l(\gamma) = 1$. Furthermore, when Γ acts faithfully, the identity of Γ is the only group element with $l(\gamma) = 0$.

We will say that a cocycle is constant if it is in $(\wedge^2 V)^\Gamma \oplus (\bigoplus_{\gamma \in \Gamma, l(\gamma)=2} \wedge^2 N^\gamma)^\Gamma$. Similarly, we will say that a cocycle is linear if it is in $(V^* \otimes \wedge^2 V)^\Gamma \oplus (\bigoplus_{\gamma \in \Gamma, l(\gamma)=2} V^{\gamma^*} \otimes \wedge^2 N^\gamma)^\Gamma$.

2.2. The Braverman-Gaitsgory conditions for PBW. Let $T(V^*) \rtimes \Gamma$ be the free $\mathbb{R}\Gamma$ -algebra generated by the bimodule $V^* \rtimes \Gamma$, and A be its quotient by the relations :

$$x \otimes y - y \otimes x - \sum_{\gamma} \pi_{\gamma}(x, y) \gamma - \sum_{\gamma} b_{\gamma}(x, y) \gamma$$

where π and b are Γ -invariant elements in $\oplus_{\gamma \in \Gamma} V^* \otimes \wedge^2(V)$ and $\oplus_{\gamma \in \Gamma} \wedge^2 V$. As before, π and b split into sums of γ -components, and the Γ -invariance is expressed in the same way as in (2). The algebra A is clearly filtered by the length of words. Following Braverman and Gaitsgory [3], the associated graded algebra $Gr(A)$ is isomorphic to $S(V^*) \rtimes \Gamma$ if and only if :

$$(4) \quad \partial_{\gamma}(\pi_{\gamma}) = 0,$$

$$(5) \quad \sum_{\alpha\beta=\gamma} \pi_{\alpha}(\pi_{\beta}(x, y), z + {}^{\beta}z) + \pi_{\alpha}(\pi_{\beta}(y, z), x + {}^{\beta}x) + \pi_{\alpha}(\pi_{\beta}(z, x), y + {}^{\beta}y) = \partial_{\gamma}(b_{\gamma}),$$

$$(6) \quad \sum_{\alpha\beta=\gamma} b_{\alpha}(\pi_{\beta}(x, y), z + {}^{\beta}z) + b_{\alpha}(\pi_{\beta}(y, z), x + {}^{\beta}x) + b_{\alpha}(\pi_{\beta}(z, x), y + {}^{\beta}y) = 0$$

for all γ in Γ and x, y, z in V^* .

When the three conditions above are satisfied, the algebra A gives a quantization of the algebra $S(V^*) \rtimes \Gamma$ for the same reason as is explained in [9](Proposition 4.5). This will be our method to obtain the quantization results of the next section.

For our purpose, let us denote $[\pi, \pi]_{\gamma} \in V^* \otimes \wedge^3 V$ and $[[b, \pi]]_{\gamma} \in \wedge^3 V$ defined by :

$$[\pi, \pi]_{\gamma}(x, y, z) := \sum_{\alpha\beta=\gamma} \pi_{\alpha}(\pi_{\beta}(x, y), z + {}^{\beta}z) + \pi_{\alpha}(\pi_{\beta}(y, z), x + {}^{\beta}x) + \pi_{\alpha}(\pi_{\beta}(z, x), y + {}^{\beta}y),$$

$$[[b, \pi]]_{\gamma}(x, y, z) := \sum_{\alpha\beta=\gamma} b_{\alpha}(\pi_{\beta}(x, y), z + {}^{\beta}z) + b_{\alpha}(\pi_{\beta}(y, z), x + {}^{\beta}x) + b_{\alpha}(\pi_{\beta}(z, x), y + {}^{\beta}y)$$

for all x, y, z in V^* .

2.3. The Gerstenhaber bracket on $H^{\bullet}(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma)$ and Poisson structures. The Gerstenhaber bracket on $H^{\bullet}(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma)$ was explicitly calculated by the first and last authors in [9]. We only recall here the results in the cases we will need and refer to [9] for a complete description.

Firstly, the Gerstenhaber bracket of two constant cocycles is zero.

Secondly, let b be a constant cocycle and π be a linear cocycle of $H^2(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma)$. Let pr_{γ} be the projection from $S(V^*) \otimes \wedge^{\bullet} V$ onto $S(V^{\gamma*}) \otimes \wedge^{\bullet-l(\gamma)} V \otimes \wedge^{l(\gamma)} N^{\gamma}$. Then the γ -component of their Gerstenhaber bracket is :

$$(7) \quad [b, \pi]_{\gamma} = \text{pr}_{\gamma} \circ [[b, \pi]]_{\gamma}.$$

Moreover, the γ -component of the Gerstenhaber bracket of π with itself is obtained by

$$(8) \quad [\pi, \pi]_{\gamma} = \text{pr}_{\gamma} \circ [[\pi, \pi]]_{\gamma}.$$

Let α and β be two elements of Γ , f_{α} be an element of $(S(V^{\alpha*}) \otimes \wedge^{\bullet-l(\alpha)} V^{\alpha} \otimes \wedge^{l(\alpha)} N^{\alpha})^{Z(\alpha)}$ and g_{β} be an element of $(S(V^{\beta*}) \otimes \wedge^{\bullet-l(\beta)} V^{\beta} \otimes \wedge^{l(\beta)} N^{\beta})^{Z(\beta)}$. If $l(\gamma) \neq l(\alpha) + l(\beta)$, then the γ component of $[f_{\alpha}, g_{\beta}]$ vanishes. Suppose that any elements in $\langle \alpha \rangle$ commutes with any elements in $\langle \beta \rangle$, where $\langle \alpha \rangle, \langle \beta \rangle$ are subsets of Γ of elements conjugate to α and β . Then the Gerstenhaber bracket of f_{α} and g_{β} is :

$$(9) \quad [f_{\alpha}, g_{\beta}]_{\gamma} = \sum_{\substack{\alpha' \in \langle \alpha \rangle, \beta' \in \langle \beta \rangle, \alpha'\beta' = \gamma \\ l(\gamma) = l(\alpha') + l(\beta')}} \text{pr}_{\gamma} \circ \{f_{\alpha'}, g_{\beta'}\},$$

where $\{f_{\alpha'}, g_{\beta'}\}$ is the usual Schouten-Nijenhuis bracket of $f_{\alpha'}$ and $g_{\beta'}$.

The first and third author defined that a Poisson structure Π on $S(V^*) \rtimes \Gamma$ is a sum of elements like

$$\pi_{\gamma} \in \left(S(V^{\gamma*}) \otimes \wedge^{2-l(\gamma)} V^{\gamma} \otimes \wedge^{l(\gamma)} N^{\gamma} \right)^{Z(\gamma)}, \quad \gamma \in C(\Gamma),$$

with $l(\gamma) = 0, 2$ satisfying $[\Pi, \Pi] = 0$.

3. PBW PROPERTY FOR CONSTANT AND LINEAR POISSON STRUCTURES

In this section, we prove that constant and linear Poisson structures (with a mild assumption in linear cases) on $S(V^*) \rtimes \Gamma$ can be quantized.

3.1. Quantization of constant Poisson structures. Following Equation (3), a Poisson bracket π on $S(V^*) \rtimes \Gamma$ splits into a sum of π_γ ,

$$\pi_0 + \sum_{\gamma} \pi_\gamma \in (S(V^*) \otimes \wedge^2 V)^\Gamma \oplus \left(\bigoplus_{\gamma \in \Gamma, l(\gamma)=2} S(V^{\gamma*}) \otimes \wedge^2 N^\gamma \right)^\Gamma.$$

We say that $\pi_0 + \sum_{\gamma} \pi_\gamma$ is a constant Poisson structure if $\pi_0 \in \wedge^2 V$ and $\pi_\gamma \in \wedge^2 N^\gamma$. We notice that in this constant case the Braverman-Gaitsgory conditions 2.2 reduce to only one condition (5), which means that π_γ has to be a cocycle. But this is automatically satisfied as we know from Equation (3) that an element in $(S(V^*) \otimes \wedge^2 V)^\Gamma \oplus \left(\bigoplus_{\gamma \in \Gamma, l(\gamma)=2} S(V^{\gamma*}) \otimes \wedge^2 N^\gamma \right)^\Gamma$ is closed with respect to the differential b_γ and $b_0 = 0$.

Theorem 3.1. *Any constant Poisson structure of $S(V^*) \rtimes \Gamma$ is quantizable.*

Proof. According to the above explanation, we know that any constant Poisson structure satisfies the Braverman-Gaitsgory conditions (4)-(6). This implies that the quotient algebra

$$H_\pi := T(V^*) \rtimes \Gamma[[\hbar]] / \langle x \otimes y - y \otimes x - \hbar(\pi_0(x, y) + \sum_{\gamma, l(\gamma)=2} \pi_\gamma(x, y)\gamma) \rangle$$

has PBW property, which defines a deformation quantization of the algebra $S(V^*) \rtimes \Gamma[[\hbar]]$ with respect to the Poisson structure $\pi = \pi_0 + \sum_{\gamma \in \Gamma, l(\gamma)=2} \pi_\gamma$. $\square$

We remark that the PBW property of the algebra H_π is checked in Etingof-Ginzburg [6]. Our proof is evident by using the results from [13].

3.2. Quantization of linear Poisson structures-abelian case. In this subsection, we will assume that Γ is an abelian group which acts faithfully on V . According to representation theory of a finite abelian group, V is decomposed into a direct sum of 1 or 2 real dimensional subspaces where Γ acts irreducibly. γ acts on 1 dimensional subspace with eigenvalue 1 or -1, and on 2 dimensional subspace by rotation of finite order.

Let π be a Poisson structure on $S(V^*) \rtimes \Gamma$, and denote π_0 its identity component. The γ -components of π are null whenever $l(\gamma) \neq 0, 2$ and take values in V^γ . As we have remarked that if $l(\gamma) = 1$, then γ action on N^γ has eigenvalue -1. There will not be any nonzero element in $V^{\gamma*} \otimes V^\gamma \otimes N^\gamma$ invariant under γ . Therefore, π_γ is possible nonzero only when $l(\gamma) = 2$ or 0. And the eigenvalues of γ action on N^γ are either -1 with multiplicity 2 or roots of unity.

Lemma 3.2. *The identity component π_0 of π defines a Lie bracket on V .*

Proof. It is a consequence of the fact that π_γ takes values in V^γ which is in the kernel of $\pi_{\gamma^{-1}}$. Therefore, the condition that the Gerstenhaber bracket $[\pi, \pi]$ vanishes at identity reduces to the Jacobi identity of π_0 . $\square$

Lemma 3.3. *Let α and β be elements of Γ with $l(\alpha) = l(\beta) = 2$. Then, $[\pi_\alpha, \pi_\beta] = 0$.*

Proof. Let x and y be the coordinates on N^α . It follows from the Γ -invariance for Poisson structure that π_α has to be β -invariant as α commutes with β for any $\beta \in \Gamma$:

$$(10) \quad \pi_\alpha({}^\beta x, {}^\beta y) = {}^\beta \pi_\alpha(x, y).$$

We observe β preserves V^α and N^α as β commutes with α . If $l(\beta) = 2$, there are three possibilities, 1) $N^\alpha \cap N^\beta = \{0\}$, 2) $\dim(N^\alpha \cap N^\beta) = 1$, 3) $N^\alpha = N^\beta$.

If $N^\alpha \cap N^\beta = \{0\}$, then x, y are β invariant for x, y in $N^{\alpha*}$. Hence by Equation (10), $\pi_\alpha(x, y) = {}^\beta\pi_\alpha(x, y)$, which shows $\pi_\alpha(x, y)$ is β invariant. Similarly, we know that π_β takes value in V^α . This shows that $[\pi_\alpha, \pi_\beta] = 0$.

If $\dim(N^\alpha \cap N^\beta) = 1$, then we know that both α and β preserves $N^{\alpha, \beta} := N^\alpha + N^\beta$ which is of 3 dimension. Furthermore, we conclude that β 's (α 's) action on N^α (on N^β) has eigenvalue 1 and -1. Hence, $N^{\alpha, \beta}$ is decomposed into a direct sum of $N_1 \oplus N_2 \oplus N_3$ such that α acts on N_1 and N_2 by -1, and N_3 by 1, and β acts on N_1 and N_3 by -1, and N_2 by 1. By Equation (10), we know that $\pi_\alpha({}^\beta x, {}^\beta y) = -\pi_\alpha(x, y) = {}^\beta\pi_\alpha(x, y)$. This shows that β acts on $\pi_\alpha(x, y)$ by -1. Similarly α acts on the image of π_β by -1. This shows that $\pi_\alpha \in N_3^* \otimes N_1 \otimes N_2$ and $\pi_\beta \in N_2^* \otimes N_1 \otimes N_3$ as N_3^* (and N_2^*) is the only 1-dim subspace of $V^{\alpha*}$ (of $V^{\beta*}$) with a nontrivial β (α) action. It is straightforward to check that $[\pi_\alpha, \pi_\beta] = 0$ in $V^* \otimes \wedge^3 V$.

If $N^\alpha = N^\beta$, β acts on $N^\alpha = N^\beta$ with determinant 1. This shows that $\pi_\alpha({}^\beta x, {}^\beta y) = \pi_\alpha(x, y)$. By Equation (10), we see that π_α takes value in V^β , and similarly π_β takes value in V^α . Direct computation shows that $[\pi_\alpha, \pi_\beta] = 0$. □

Theorem 3.4. *Let Γ be a finite abelian group which acts faithfully on a finite dimensional vector space V . Then any linear Poisson structure π of $S(V^*) \rtimes \Gamma$ is quantizable.*

Proof. Following the above lemmas, $[\pi, \pi]_\gamma$ reduces to :

$$\begin{aligned} [\pi, \pi]_\gamma(x, y, z) &= \pi_0(\pi_\gamma(x, y), z + {}^\gamma z) + \pi_0(\pi_\gamma(y, z), x + {}^\gamma x) + \pi_0(\pi_\gamma(z, x), y + {}^\gamma y) \\ &\quad + 2 \times (\pi_\gamma(\pi_0(x, y), z) + \pi_\gamma(\pi_0(y, z), x) + \pi_\gamma(\pi_0(z, x), y)). \end{aligned}$$

This expression is zero whenever $l(\gamma) \neq 2$. Suppose now that $l(\gamma) = 2$. If x, y, z are all in V^γ , we get 0 since V^γ is the kernel of π_γ . If two of x, y, z are in V^γ , we also get 0 for the same reason and because π_0 is γ -invariant. Suppose now that x and y are in N^γ and that z is γ -invariant. Then, $[\pi, \pi]_\gamma(x, y, z)$ lies in V^γ , and it is zero from the fact that π is a Poisson bracket (Section 2.3): $[\pi, \pi]_\gamma = 0$. □

3.3. An important example. In order to prove the quantization theorem for general linear Poisson structures, we consider an important example in this subsection.

Let $\rho = \exp(\frac{2\pi i}{2n+1})$. Denote $\alpha_k = \begin{pmatrix} \rho^k & 0 \\ 0 & \rho^{-k} \end{pmatrix}$, and $\beta_k = \begin{pmatrix} 0 & \rho^{-k} \\ \rho^k & 0 \end{pmatrix}$, and $\Gamma_n = \{\alpha_k, \beta_l : 0 \leq k, l \leq 2n\}$. Γ_n is a finite group of order $4n+2$ acting faithfully on $V = \mathbb{C}^2$, a complex 2-dim and real 4-dim vector space. α_k 's eigenvalues are ρ^k and ρ^{-k} , and β_k 's eigenvalues are ± 1 . Let z_1, z_2 be complex coordinate functions on V .

We consider linear Poisson structures on $S(V^*) \rtimes \Gamma_n$. We first look at the Poisson structure on the identity component. As α_k acts on V diagonally, α_k acts on $V^* \otimes \wedge^2 V$ also diagonally with eigenvalues $\rho^{3k}, \rho^k, \rho^{-k}$, and ρ^{-3k} . If $\rho^{3k} \neq 1$, there is no non zero linear bivector field on V , which is α_k invariant. Accordingly, if $\rho^3 \neq 1$, there is no Γ_n -invariant linear Poisson structure π_0 on V . If $\rho^3 = 1$, then π_0 is a linear combination of $z_1 \bar{\partial}_1 \wedge \partial_2$, $z_2 \partial_1 \wedge \bar{\partial}_2$, and $\bar{z}_1 \partial_1 \wedge \bar{\partial}_2$, $\bar{z}_2 \bar{\partial}_1 \wedge \partial_1$. If we assume that π_0 to be real, then we have

$$\pi_0 = az_1 \bar{\partial}_1 \wedge \partial_2 + \bar{a} \bar{z}_1 \partial_1 \wedge \bar{\partial}_2 + bz_2 \partial_1 \wedge \bar{\partial}_2 + \bar{b} \bar{z}_2 \bar{\partial}_1 \wedge \partial_2.$$

Furthermore by invariance with respect to the β_k 's action, we have $a = -b$ and $\bar{a} = -\bar{b}$ in the above equation, i.e.

$$(11) \quad \pi_0 = a(z_1 \bar{\partial}_1 \wedge \partial_2 - z_2 \partial_1 \wedge \bar{\partial}_2) + \bar{a}(\bar{z}_1 \partial_1 \wedge \bar{\partial}_2 - \bar{z}_2 \bar{\partial}_1 \wedge \partial_2).$$

Observe β_k has real codimension=2 fixed point subspace, while α_k only fixes the origin of V . $V^k := V^{\beta_k}$ is determined by $\rho^k z_1 - z_2 = \rho^{-k} \bar{z}_1 - \bar{z}_2 = 0$. The normal subspace N^k to V^k is determined by $\rho^k z_1 + z_2 = \rho^{-k} \bar{z}_1 + \bar{z}_2 = 0$. Vector fields along N^k are spanned by $\rho^{-k} \partial_1 - \partial_2$ and $\rho^k \bar{\partial}_1 - \bar{\partial}_2$. Therefore, the Poisson structure at β_k component can be written as

$$\Pi_k = [c_k(\rho^k z_1 + z_2) - \bar{c}_k(\rho^{-k} \bar{z}_1 + \bar{z}_2)] (\rho^{-k} \partial_1 - \partial_2) \wedge (\rho^k \bar{\partial}_1 - \bar{\partial}_2).$$

Furthermore, as $\alpha_l \beta_k \alpha_l^{-1} = \alpha_{k-2l}$, by invariance of Π_k with respect to the conjugation action of Γ_k , $c_{2k} = c_0 \rho^{-k}$, $0 \leq k \leq 2n$. Therefore, we have

$$(12) \quad \begin{aligned} \Pi_{2k} = & [c_0(\rho^k z_1 + \rho^{-k} z_2) - \bar{c}_0(\rho^{-k} \bar{z}_1 + \rho^k \bar{z}_2)] \\ & (\partial_1 \wedge \bar{\partial}_1 - \rho^{-2k} \partial_1 \wedge \bar{\partial}_2 - \rho^{2k} \partial_2 \wedge \bar{\partial}_1 + \partial_2 \wedge \bar{\partial}_2), \quad 0 \leq k \leq 2n. \end{aligned}$$

In summary, a Poisson structure Π on $S(V^*) \rtimes \Gamma_n$ is of the form, $\Pi = \pi_0 + \sum_{k=0}^{2n} \Pi_{2k}$ where Π_{2k} is defined as in Equation (12), and π_0 vanishes unless $n = 3$. When $n = 3$, π_0 is defined as in Equation (11)

By the same reason as in the proof of Theorem 3.4, we conclude that if we assume $[\pi_0, \Pi_k] = 0$, then $[[\pi_0, \Pi_k] + [\Pi_k, \pi_0]] = 0$ for any k .

From Equation (5), we see that as $\dim(N^\beta) = 2$ the $[[\pi_\alpha, \pi_\beta]](x, y, z) = 0$ if x, y, z are all along the normal direction N^β . Furthermore, as π_β is a multiple of the highest wedge power of the normal direction N^β , to have non-zero outcome two of the three x, y, z have to be from the normal direction N^β . This implies that $[[\pi_\alpha, \pi_\beta]]$ as an element in $V^* \otimes \wedge^3 V$ is equal to the Schouten-Nijenhuis bracket $[\pi_\alpha, \pi_\beta]$.

We use this observation to compute $[[\Pi_{2k}, \Pi_{2l}]]$. A long but straightforward computation leads to the following result at α_{2k-2l} ,

$$\begin{aligned} [[\Pi_{2k}, \Pi_{2l}]] = & [c_0(\rho^k z_1 + \rho^{-k} z_2) - \bar{c}_0(\rho^{-k} \bar{z}_1 + \rho^k \bar{z}_2)] \\ & \times \left[(2\bar{c}_0 \rho^l - \bar{c}_0 \rho^{-l+2k} - \bar{c}_0 \rho^{3l-2k}) \partial_1 \wedge \partial_2 \wedge \bar{\partial}_1 \right. \\ & + (-2\bar{c}_0 \rho^{-l} + \bar{c}_0 \rho^{l-2k} + \bar{c}_0 \rho^{-3l+2k}) \partial_1 \wedge \partial_2 \wedge \bar{\partial}_2 \\ & + (2c_0 \rho^{-l} - c_0 \rho^{l-2k} - c_0 \rho^{-3l+2k}) \bar{\partial}_1 \wedge \partial_1 \wedge \bar{\partial}_2 \\ & \left. + (-2c_0 \rho^l + c_0 \rho^{-l+2k} + c_0 \rho^{3l-2k}) \bar{\partial}_1 \wedge \partial_2 \wedge \bar{\partial}_2 \right]. \end{aligned}$$

Define for $0 \leq k \leq 2n$,

$$(13) \quad B_{2k} := (2n+1)(\rho^k - \rho^{-k}) [-|c_0|^2 \partial_2 \wedge \bar{\partial}_2 + |c_0|^2 \partial_1 \wedge \bar{\partial}_1 + (\bar{c}_0)^2 \partial_1 \wedge \partial_2 - c_0^2 \bar{\partial}_1 \wedge \bar{\partial}_2].$$

With a long but straightforward computation, we are able to prove

$$\sum_{p-q=2k, 0 \leq q \leq 2n} [[\Pi_{2p}, \Pi_{2q}]] = \partial^{\alpha_{2k}} B_{2k}.$$

And it is not difficult to compute that

$$(14) \quad [[B_{2k}, \Pi_{2l}]] = 0, \quad 0 \leq k, l \leq 2n;$$

and

$$[[B_{2k}, \pi_0]] = 0, \quad 0 \leq k \leq 2n, \quad n = 3.$$

Therefore, we conclude with the following proposition

Proposition 3.5. *For Γ_n action on $V = \mathbb{C}^2$, any linear Poisson structures on $S(V^*) \rtimes \Gamma_n$ can be quantized.*

Proof. By the above computation, we see that the relation defining

$$H_\Pi := T(V) \rtimes \Gamma_n[[\hbar]]$$

$$\left\langle x \otimes y - y \otimes x - \hbar \left(\pi_0(x, y) + \sum_{0 \leq k \leq 2n} \Pi_{2k}(x, y) \beta_{2k} \right) - \hbar^2 \left(\sum_{1 \leq k \leq 2n} B_{2k}(x, y) \alpha_{2k} \right) \right\rangle$$

satisfies the Braverman-Gaitsory conditions (4)-(6). This implies that H_Π has PBW property, which shows that H_Π is a deformation quantization of $S(V^*) \rtimes \Gamma_n$ along the direction defined by Π . $\square$

Remark 3.6. *We point out that in the proof of Proposition 3.5, there have to be nonzero terms B_{2k} for $0 \leq k \leq 2n$ as $[\pi, \pi]$ is not zero. This is different from the standard PBW theorem for Lie algebras where B_{2k} can be chosen to be zero. We will see in the following subsection that this example is essentially the only case that B_{2k} has to be nonzero.*

3.4. Quantization of linear Poisson structures-general case. In this subsection Γ is a finite group (not necessary abelian) acting faithfully on a vector space V . We assume that V is equipped with a Γ -invariant complex structure. We prove the following theorem.

Theorem 3.7. *Let Γ be a finite group acting faithfully on a complex vector space V . Any real linear Poisson structure on $S(V^*) \rtimes \Gamma$ is quantizable.*

The proof of this theorem consists of several steps. We start with recalling some results about finite subgroups of $GL(2, \mathbb{C})$.

Lemma 3.8. *A nonabelian finite subgroup G of $SL(2, \mathbb{C})$ must contain the element $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \in SL(2, \mathbb{C})$.*

Proof. We notice that the canonical action of G on $\mathbb{C}^2$ is irreducible. Otherwise, G will be a subgroup of $GL(1, \mathbb{C}) \times GL(1, \mathbb{C})$, which is abelian. Therefore, according to [14][Chapter 6, Proposition 17], the order of the center of G is divisible by 2. Therefore, there is an element in G of order 2. As $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ is the unique element in $SL(2, \mathbb{C})$ of order 2, we conclude that if G is not abelian, G contains $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$. $\square$

The following lemma is a corollary of [5][§26, Theorem 26].

Lemma 3.9. *Let Γ be a nonabelian finite subgroup of $GL(2, \mathbb{C})$. If Γ does not contain any matrix of the form $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ for $a \neq 1$, then there is a natural number n such that Γ is conjugate to the group Γ_n as is introduced in subsection 3.3.*

Proof. We start by considering the intersection $G = \Gamma \cap SL(2, \mathbb{C})$. If G is trivial, then there is an injective group homomorphism

$$\Gamma \rightarrow GL(2, \mathbb{C})/SL(2, \mathbb{C}) \cong \mathbb{C} - \{0\}.$$

This shows that Γ is abelian, which contradicts the assumption that Γ is not abelian. Therefore, G is a nontrivial subgroup of $SL(2, \mathbb{C})$. The following discussion is divided into two parts according to whether G is abelian.

- G is not abelian. Then by Lemma 3.8, G contains the element $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$, which is in the center of G . This contradicts to the assumption of this lemma.
- G is abelian. By conjugation with an invertible matrix, we can assume that G contains a diagonal element like $A = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$ with $a \neq -1$. (We remark that any element in Γ is diagonalizable as Γ is of finite order.) Furthermore, we recall the fact that if $B = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ commutes with A , then $\beta = \gamma = 0$. Hence, any element in G is of the form $B = \begin{pmatrix} \beta & 0 \\ 0 & \beta^{-1} \end{pmatrix}$ for $\beta \in \mathbb{C} - \{0\}$. Therefore, we conclude that G is a cyclic subgroup of $SL(2, \mathbb{C})$ isomorphic to $\left\{ \begin{pmatrix} \rho & 0 \\ 0 & \rho^{-1} \end{pmatrix} : \rho^{2n+1} = 1 \right\}$ for some $n \in \mathbb{N}$. (If $\rho^{2n} = 1$, then $\rho^n = -1$ and G contains the element $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$.)

We observe that if $B = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in GL(2, \mathbb{C})$ is a normalizer of G , then $\alpha\beta = \gamma\delta = 0$. Therefore, $B = \begin{pmatrix} \alpha & 0 \\ 0 & \gamma \end{pmatrix}$ or $\begin{pmatrix} 0 & \beta \\ \delta & 0 \end{pmatrix}$. This shows that any element in Γ is either diagonal or of the form $\begin{pmatrix} 0 & \beta \\ \delta & 0 \end{pmatrix}$. As we have assumed that Γ is not an abelian group, there has to be a nonzero B in Γ of the form $\begin{pmatrix} 0 & \beta \\ \delta & 0 \end{pmatrix}$.

Compute $B^2 = \begin{pmatrix} \beta\delta & 0 \\ 0 & \delta\beta \end{pmatrix}$. By the assumption of Γ , $\beta\delta = 1$. Therefore, $B = \begin{pmatrix} 0 & a \\ a^{-1} & 0 \end{pmatrix} \in \Gamma$. Now choose $U = \begin{pmatrix} 0 & a^{\frac{1}{2}} \\ a^{-\frac{1}{2}} & 0 \end{pmatrix}$, and consider the group $\tilde{\Gamma} = U^{-1}\Gamma U$ which is again not abelian and does not contain any matrix of the form $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$. Under this isomorphism, we see that $\tilde{G} = G = \left\{ \begin{pmatrix} \rho & 0 \\ 0 & \rho^{-1} \end{pmatrix} : \rho^{2n+1} = 0 \right\}$ and $\tilde{\Gamma}$ contains a matrix $\beta_0 = U^{-1}BU = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \in \tilde{\Gamma}$.

Now if there is any other element C in $\tilde{\Gamma}$ of the form $\begin{pmatrix} 0 & \beta' \\ \delta' & 0 \end{pmatrix}$ then by the same arguments as B , we know that $\beta' = 1/\delta'$. Furthermore, as $\beta_0 B = \begin{pmatrix} \delta' & 0 \\ 0 & \delta'^{-1} \end{pmatrix} \in \tilde{G} = G$. This implies that $B = \begin{pmatrix} 0 & \rho \\ \rho^{-1} & 0 \end{pmatrix}$ with $\rho^{2n+1} = 1$.

Next if there is any element $D = \begin{pmatrix} \alpha & 0 \\ 0 & \gamma \end{pmatrix}$ in $\tilde{\Gamma}$, compute $\beta_0 D \beta_0 D = \begin{pmatrix} \alpha\gamma & 0 \\ 0 & \alpha\gamma \end{pmatrix}$. As $\tilde{\Gamma}$ has no element like $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ with $a \neq 1$, we conclude that $\alpha = 1/\gamma$ and D belongs to G .

Summarizing the above analysis, we have seen that there exists $n \in \mathbb{N}$, such that $\tilde{\Gamma} = \left\{ \begin{pmatrix} 0 & \rho^i \\ \rho^{-i} & 0 \end{pmatrix}, \begin{pmatrix} \rho^i & 0 \\ 0 & \rho^{-i} \end{pmatrix} : 0 \leq i \leq 2n, \rho^{2n+1} = 1 \right\}$.

□

Proof of theorem 3.7:

With Theorem 3.4, it is sufficient to work with Γ which is nonabelian. We choose a Γ -invariant hermitian metric on V which always exists as Γ is finite. We prove that there exists a choice for B_γ such that the Braverman-Gaitsory conditions (4)-(6) are satisfied.

Step I: The Braverman-Gaitsory condition (4) is satisfied automatically by the assumption on π_α .

Step II: In the following, we make a proper choice for B_γ with $l(\gamma) = 4$ such that the Braverman-Gaitsory condition (5) is satisfied.

As Γ is acting on a complex vector space, the fixed subspace of any group element γ is of even real codimension. Let π_0 be the linear Poisson structure at the identity component, π_α be the linear Poisson structure at the α component with $l(\alpha) = 2$.

We look at $[\pi_0, \pi_\alpha]$, $[\pi_\alpha, \pi_0]$, and $[\pi_\alpha, \pi_\beta]$ with $l(\alpha) = l(\beta) = 2$. By the same arguments as in the proof of Theorem 3.4, we have that for any α with $l(\alpha) = 2$, $[\pi_0, \pi_\alpha] + [\pi_\alpha, \pi_0] = 0$ as $\pi_0 + \sum_\alpha \pi_\alpha$ is a Poisson structure. Therefore, we are reduced to look at $[\pi_\alpha, \pi_\beta]$.

We observe that if $V^\alpha = V^\beta$, then $\wedge^2 N^\alpha = \wedge^2 N^\beta$ vanishes on functions depending only on variables in $V^\alpha = V^\beta$. It is easy to compute that $[\pi_\alpha, \pi_\beta] = 0$. This reduces us to the situation that $V^\alpha \neq V^\beta$.

When $V^\alpha \neq V^\beta$, we have the equation $V^\alpha + V^\beta = V$ as both V^α and V^β are complex subspaces of V of complex codimension 1. Furthermore the equality $V^\alpha + V^\beta = V$ implies that $V^{\alpha\beta} = V^\alpha \cap V^\beta$ which is of complex codimension 2 and real codimension 4. We notice that in this case both α and β fix every point in $V^{\alpha\beta}$, and also preserve the normal direction $N^{\alpha\beta}$. We are interested in $[\pi_\alpha, \pi_\beta]$, which is now at $\alpha\beta$ component. As $[\pi_\alpha, \pi_\beta]$ is only a tri-vector field supported at $N^{\alpha\beta}$ with $l(\alpha\beta) = 4$, we know that $[\pi_\alpha, \pi_\beta]$ is 0 in the Hochschild cohomology of $S(V) \rtimes \Gamma$.

Now we fix an element γ with $l(\gamma) = 4$, then we know from the previous paragraph that if $[\pi_\alpha, \pi_\beta]$ is nonzero at $\gamma = \alpha\beta$, then α and β acts on V preserving N^γ and fixing every element in V^γ . This leads us to look at the subgroup Γ_γ of Γ whose elements act trivially on V^γ . Γ_γ contains all α such that $V^\gamma \subset V^\alpha$, which acts on N^γ faithfully.

Let $\alpha \in \Gamma_\gamma$ with $l(\alpha) = 2$. By the assumption on π_α , it is an element in $V^{\alpha*} \otimes \wedge^2 N^\alpha$, which can be written as a sum of two terms $\pi_\alpha^1 + \pi_\alpha^2$ as $V^{\alpha*} = V^{\gamma*} \oplus (N^{\gamma*} \cap V^{\alpha*})$. We easily see that π_α^1 will

not contribute to $[\pi_\alpha, \pi_\beta]$ as N^α and N^β are orthogonal to V^γ . This shows that to study $[\pi_\alpha, \pi_\beta]$, it is enough to assume that π_α belongs to $N^{\gamma*} \otimes \wedge^2 N^\alpha$. Furthermore, if n_α is the holomorphic vector along N^α and v_α is the holomorphic vector along $V^{\alpha*}$ in N^γ , then we can write $\pi_\alpha = (c_\alpha v_\alpha - \bar{c}_\alpha \bar{v}_\alpha) n_\alpha \wedge \bar{n}_\alpha$ for some complex number c_α by the fact that π_α is real.

If Γ_γ is abelian, then by Lemma 3.3, we know that $[\pi_\alpha, \pi_\beta] = 0$ for any α, β , and therefore we set $B_\delta = 0$ for $\delta \in \Gamma_\gamma$. In the following, we assume that Γ_γ is not abelian. If Γ_γ contains an element ν which acts on N^γ of the form $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ with a unitary number $a \neq 1$, then it is easy to check $\nu_*(\pi_\alpha) = (a^{-1}c_\alpha v_\alpha - \bar{a}^{-1}\bar{c}_\alpha \bar{v}_\alpha) n_\alpha \wedge \bar{n}_\alpha$ is not invariant under ν unless $c_\alpha = 0$. This shows that π_α has to be zero if it is invariant under ν and therefore $[\pi_\alpha, \pi_\beta] = 0$ in this case. We choose $B_\gamma = 0$ for this type of γ . Therefore, for nonzero B_γ , we only need to consider the situation that Γ_γ is not abelian and contains no element of the form $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ with $a \neq 1$. By Lemma 3.9, Γ_γ action on N^γ is isomorphic to the situation studied in subsection 3.3. And we can choose B_γ as Equation (13).

Step III: we prove that $\sum_{\alpha, \gamma, l(\alpha) \leq 2, l(\gamma)=4} [B_\gamma, \pi_\alpha] = 0$ with the choices of B_γ introduced in Step II.

We decompose the above sum into 2 parts

- (1) $\sum_{\alpha, \gamma, l(\alpha) \leq 2, l(\gamma)=4, l(\gamma\alpha) \geq 4} [B_\gamma, \pi_\alpha];$
- (2) $\sum_{\alpha, \gamma, l(\alpha) \leq 2, l(\gamma)=4, l(\gamma\alpha)=2} [B_\gamma, \pi_\alpha].$

We consider the part of sum with $l(\gamma\alpha) \geq 4$. Let $\delta = \gamma\alpha$. Define $D_\delta = \sum_{\gamma\alpha=\delta, l(\alpha) \leq 2, l(\gamma)=4} [B_\gamma, \pi_\alpha]$, which is an element in $\wedge^3 V$. We notice that D_δ is ∂^δ closed as

$$\begin{aligned} \partial(\sum_{\alpha, \gamma} [B_\gamma, \pi_\alpha]) &= \sum_{\alpha, \gamma} [\partial B_\gamma, \pi_\alpha] \\ &= \sum_{\alpha, \gamma=\beta\lambda} [[\pi_\beta, \pi_\lambda], \pi_\alpha] \\ &= 0. \end{aligned}$$

According to Equation (3), we see that D_γ in $\wedge^3 V$ is a zero cocycle as $l(\gamma) \geq 4$. On the other hand, we see that D_γ is in $\wedge^3 V$. If we define elements in V of degree -1 and elements in V^* of degree 1, then D_γ is of degree -3, and ∂^γ is of degree 0. We see that $S(V^*) \otimes \wedge^2 V$ has degree greater or equal to -2. Therefore, $\partial^\gamma(S(V^*) \otimes \wedge^2 V)$ has no term with degree less than -2 since $\deg(\partial^\gamma) = 0$. This shows that if $D_\gamma \neq 0$, it cannot be a coboundary of ∂^γ . This shows that D_γ has to be zero as D_γ is a trivial cocycle by Equation (3).

We consider the part of sum with $l(\gamma\alpha) \leq 2$. This implies that $V^\gamma \subset V^\alpha$ because otherwise $V^\gamma + V^\alpha = V$ and $V^{\gamma\alpha} = V^\gamma \cap V^\alpha$ which is of real codimension 6. In this case, we know that $\gamma\alpha$ is also in Γ_γ . Therefore, we can use Equation (14) to conclude that $[B_\gamma, \pi_\alpha] = 0$.

In conclusion, Steps I-III show that Braverman-Gaiety conditions (4)-(6) are satisfied for any linear Poisson structures on $S(V^*) \rtimes \Gamma$ and a proper choice of B_γ . Therefore, by PBW property, we see that the algebra

$$H_\Pi := T(V) \rtimes \Gamma[[\hbar]]$$

$$\left\langle x \otimes y - y \otimes x - \hbar(\pi_0(x, y) + \sum_{\alpha} \pi_\alpha(x, y)\alpha) - \hbar^2(\sum_{\gamma} B_\gamma(x, y)\gamma) \right\rangle$$

defines a deformation quantization of $S(V^*) \rtimes \Gamma$ along the direction of $\pi_0 + \sum_{\alpha} \pi_\alpha$. $\square$

4. HOCHSCHILD COHOMOLOGY AND NON COMMUTATIVE POISSON COHOMOLOGY

In this section, we would like to study various properties and examples of the Poisson structures and algebras we constructed in the previous section. To state our results, we fix some convention. Note that if α and β are conjugate to each other inside Γ , then $l(\alpha) = l(\beta)$. Therefore, it is legitimate to define **codimension** of a conjugacy class of Γ by the codimension of an element $\alpha \in \Gamma$. Define c_k to be the

number of conjugacy class of Γ with codimension k . In the following, we use $\mathbb{R}((\hbar))$ to stand for the algebra of Laurent Polynomials of $\hbar$. For any algebra A over the ring $\mathbb{R}[[\hbar]]$, we use $A((\hbar))$ to stand for the extension $A \otimes_{\mathbb{R}[[\hbar]]} \mathbb{R}((\hbar))$.

Proposition 4.1. *The periodic cyclic homology of the algebra H_π for any constant or linear Poisson structures on $S(V^*) \rtimes \Gamma$ is equal to*

$$\begin{aligned} HP_0(H_\pi((\hbar))) &= \sum_k \mathbb{R}((\hbar))^{\times c_k} = \mathbb{R}((\hbar))^{\times |C(\Gamma)|} \\ HP_1(H_\pi((\hbar))) &= 0. \end{aligned}$$

Proof. By Getzler-Goodwillie [7], we know that as the periodic cyclic homology is invariant under deformation, $HP_\bullet(H_\pi((\hbar)))$ is equal to $HP_\bullet(S(V^*) \rtimes C(\Gamma))((\hbar))$. By the computation in [4], $HP_\bullet(S(V^*) \rtimes \Gamma)$ is equal as is stated above. $\square$

4.1. Hochschild cohomology of symplectic reflection algebra. In this subsection, we restrict ourselves to quantization of a special type of constant Poisson structures. These are called symplectic reflection algebras. Let V be a symplectic vector space with standard symplectic 2-form ω . Let π be the associated Poisson structure of ω . For γ with $l(\gamma) = 2$, consider the restriction ω_γ of ω to N^γ . ω_γ is an invertible bilinear operation on N^γ . Define π_γ be the inverse of ω_γ . We choose $c_\gamma \in \mathbb{R}$ for all γ with $l(\gamma) = 2$ with $c_{\alpha\gamma\alpha^{-1}} = c_\gamma$ for all $\alpha, \gamma \in \Gamma$. Define a constant Poisson structure Π on $S(V^*) \rtimes \Gamma$ by

$$\Pi = \pi + \sum_{\gamma, l(\gamma)=2} c_\gamma \pi_\gamma.$$

By Thm 3.1, $S(V^*) \rtimes \Gamma$ has a deformation quantization with respect to Π . This algebra can be written as

$$H_{\omega,c} = T(V^*) \otimes \Gamma[[\hbar]] / \langle x \otimes y - y \otimes x - \hbar(\pi(x, y) + \sum_{\gamma, l(\gamma)=2} c_\gamma \pi_\gamma(x, y) \gamma) \rangle.$$

This algebra is called symplectic reflection algebra by Etingof and Ginzburg [6].

To compute the Hochschild cohomology of $H_{\omega,c}$, we compute the Poisson cohomology of Π first. We recall that the Poisson cohomology of a Poisson structure Π on an algebra A is defined to be the cohomology of the complex $H^\bullet(A, A)$ with the differential $\partial(a) = [\Pi, a]$, where $[\ , \]$ is the Gerstenhaber bracket on $H^\bullet(A, A)$.

Proposition 4.2. *The Poisson cohomology $H_\Pi^\bullet(S(V^*) \rtimes \Gamma)$ is equal to*

$$H_\Pi^\bullet(S(V^*) \rtimes \Gamma) = \mathbb{R}^{\times c_\bullet}.$$

Proof. We introduce a grading on

$$H^\bullet(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma) = \left(\bigoplus_{\gamma \in \Gamma} S(V^{\gamma*}) \otimes \wedge^{\bullet-l(\gamma)} V^\gamma \otimes \wedge^{l(\gamma)} N^\gamma \right)^\Gamma$$

by setting elements in $S(V^{\gamma*}) \otimes \wedge^{\bullet-l(\gamma)} V^\gamma \otimes \wedge^{l(\gamma)} N^\gamma$ of degree $l(\gamma)$.

By Poisson cohomology with respect to Π , we mean the cohomology on $H^\bullet(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma)$ with the differential d_Π defined by taking the generalized Schouten-Nijenhuis bracket $d_\Pi f = [\Pi, f]$ for $f \in H^\bullet(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma)$. According to [9][Thm. 3.4], the Poisson differential d_Π is compatible with the above filtration with respect to $l(\gamma)$ as if $\deg(f) = i$, $(d_\Pi f)_\gamma = 0$ if $l(\gamma) \neq i$ or $i + 2$. Therefore, we can use spectral sequence associated to the filtration defined by the grading l to compute the Poisson cohomology H_Π .

The E_0 of the spectral sequence associated to the above filtration is the Poisson cohomology with respect to the Poisson structure π which is the component of Π supported at identity of the group Γ on the graded complex $\text{Gr}(H^\bullet(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma))$, i.e.

$$\text{Gr}^p(H^\bullet(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma)) = \left(\bigoplus_{\gamma, l(\gamma)=p} S(V^{\gamma*}) \otimes \wedge^{\bullet-l(\gamma)} V^\gamma \otimes \wedge^{l(\gamma)} N^\gamma \right)^\Gamma.$$

As ω is a symplectic form, the Poisson cohomology of π can be computed easily

$$E_1^{p,q} = \begin{cases} H_\pi^p(S(V^*) \rtimes \Gamma) = \mathbb{R}^{\times c_p}, & q = 0, \\ \{0\}, & q \neq 0. \end{cases}$$

We notice that the codimension of any group element γ is even because γ preserves the symplectic structure. Therefore, $E_1^{p,q} = 0$ if and only if $p + q$ is odd. This implies that the spectral sequence degenerates at E_1 . Therefore we have

$$H_\Pi^\bullet(S(V^*) \rtimes \Gamma) = H_\pi^\bullet(S(V^*) \rtimes \Gamma) = \mathbb{R}^{\times c_\bullet}.$$

□

Theorem 4.3. *The Hochschild cohomology of $H_{\omega,c}((\hbar))$ is computed as follows,*

$$H^\bullet(H_{\omega,c}((\hbar)), H_{\omega,c}((\hbar))) = \mathbb{R}((\hbar))^{\times c_\bullet}.$$

Proof. The proof of this theorem uses the spectral sequence with respect to the $\hbar$ -filtration. The E_0 terms are the Poisson cohomology. We notice that $H_\Pi^\bullet(S(V^*) \rtimes \Gamma)$ is trivial when $\bullet$ is odd, and conclude that the spectral sequence degenerated at E_1 . Therefore, the Hochschild (co)homology equals the Poisson (co)homology. □

Remark 4.4. *Theorem 4.3 is a generalization of [6][Theorem 1.8 (i)]. Here, with more information about the generalized Schouten-Nijenhuis bracket, we are able to avoid the restriction of [6] on “except possibly a countable set”.*

In Thm 4.3, we see that for a constant Poisson structure Π , if the Poisson structure at the identity component π_0 is the inverse of a symplectic structure, then its Poisson cohomology and Hochschild cohomology are determined by π_0 completely.

In the following we show one example that if the Poisson structure π_0 is degenerated, then the Poisson cohomology of Π depends also on the information of Π at other conjugacy classes.

Consider $(V = \mathbb{R}^2, \omega = dx \wedge dy)$, and $\mathbb{Z}_2 := \mathbb{Z}/2\mathbb{Z} = \{1, e\}$ acts on $\mathbb{R}^2$ by $e(x, y) = (-x, -y)$. Denote $\pi = -\partial_x \wedge \partial_y$. Observe that 0 is the only fixed point of e . For any constant c , consider $\Pi = c\pi e : x \wedge y \mapsto c\pi(x, y)e$ which defines a constant Poisson structure on $S(V^*) \rtimes \mathbb{Z}_2$. In the following, we show that the Poisson cohomology of Π does distinguish Π from the trivial Poisson structure and also those Poisson structures in Proposition 4.2.

Proposition 4.5.

$$\begin{aligned} H_\Pi^0 &= (S(V^*))^{\mathbb{Z}_2} \\ H_\Pi^1 &= \{f_1 \partial_x + f_2 \partial_y \in (S(V^*) \otimes V)^{\mathbb{Z}_2} : \partial_x f_1(0) + \partial_y f_2(0) = 0\} \\ H_\Pi^2 &= (S(V^*) \otimes \wedge^2 V)^{\mathbb{Z}_2}. \end{aligned}$$

Proof. The computation for H^0 are trivial as there is only degree=2 elements supported at e by Equation (3).

For H^1 , we first observe that the image of d^Π on $(S(V^*))^{\mathbb{Z}_2}$ is trivial by Equation (3). For the kernel of d^Π , we compute

$$[f_1 \partial_x + f_2 \partial_y, \Pi] = c(\partial_x(f_1) + \partial_y(f_2))\partial_x \wedge \partial_y|_0,$$

where $c(\partial_x(f_1) + \partial_y(f_2))\partial_x \wedge \partial_y|_0$ is the restriction of $c(\partial_x(f_1) + \partial_y(f_2))\partial_x \wedge \partial_y$ to the origin 0. Therefore, $H_\Pi^1 = \{f_1 \partial_x + f_2 \partial_y \in (S(V^*) \otimes V)^{\mathbb{Z}_2} : \partial_x f_1(0) + \partial_y f_2(0) = 0\}$.

For H_Π^2 , we notice that d^Π vanishes as there is no higher degree terms. For the image of d^Π , from the previous computation, we see that $\text{Im}(d^\Pi) = \mathbb{R}\partial_x \wedge \partial_y|_0$. Therefore we conclude from Equation (3) that $H_\Pi^2 = (S(V^*) \otimes \wedge^2 V)^{\mathbb{Z}_2}$. □

4.2. Hochschild cohomology of $\mathcal{U}(\mathfrak{g}) \rtimes \Gamma$. Let $V = \mathfrak{g}$ be a Lie algebra such that its bracket is Γ invariant. The Lie bracket on $\mathfrak{g}$ defines a Poisson structure on V^* , which also defines a Poisson structure on $S(V) \rtimes \Gamma$. Then, $\mathcal{U}(\mathfrak{g}) \rtimes \Gamma$ is a quantization of π on $S(V) \rtimes \Gamma$. We notice that the Poisson bracket π does not have any other γ -component for γ different from the unity, i.e. $\pi = \pi_0$.

Now, $\mathcal{U}(\mathfrak{g}) \rtimes \Gamma$ is a filtered Koszul algebra over $\mathbb{R}\Gamma$. Therefore, it has a Koszul resolution and a small complex which calculates its Hochschild cohomology. This complex splits into a direct sum of subcomplexes:

$$(CK^\bullet(\mathcal{U}(\mathfrak{g}) \rtimes \Gamma), \partial) = \bigoplus_{\gamma \in C(\Gamma)} (CK_\gamma^\bullet, \partial_\gamma)$$

where $CK_\gamma^\bullet = (\mathcal{U}(\mathfrak{g}) \otimes \wedge^\bullet \mathfrak{g}^*)^{Z(\gamma)}$ and :

$$\begin{aligned} \partial_\gamma(f)(x_0, \dots, x_n) &= \sum_{i=0}^n (-1)^i f(x_0, \dots, \widehat{x}_i \dots, x_n)(x_i -^\gamma x_i) \\ &+ \sum_{i=0}^n (-1)^i [x_i, f(x_0, \dots, \widehat{x}_i \dots, x_n)] + \sum_{i < j} (-1)^{j-i-1} f(x_0, \dots, x_{i-1}, [x_i, x_j], x_{i+1}, \dots, \widehat{x}_j \dots, x_n). \end{aligned}$$

The brackets in the above formula stand for the Lie bracket of $\mathfrak{g}$ and for the action of $\mathfrak{g}$ on $\mathcal{U}(\mathfrak{g})$. Notice that the PBW property of $\mathcal{U}(\mathfrak{g})$ implies that the symmetrization map from $S\mathfrak{g}$ to $\mathcal{U}(\mathfrak{g})$ is an isomorphism of $\mathfrak{g}$ -modules as well as Γ -modules.

According to 2.3, the Poisson complex splits as well in a direct sum of subcomplexes

$$(C_\pi^\bullet(\mathcal{U}(\mathfrak{g}) \rtimes \Gamma), \partial^\pi) = \bigoplus_{\gamma \in C(\Gamma)} (C_\gamma^\bullet, \partial_\gamma^\pi),$$

where $C_\gamma^\bullet = (S(\mathfrak{g}^\gamma) \otimes \wedge^{\bullet-l(\gamma)} \mathfrak{g}^{\gamma*} \otimes \wedge^{l(\gamma)} N^{\gamma*})^{Z(\gamma)}$ with the differential :

$$\begin{aligned} \partial_\gamma^\pi(f)(x_0, \dots, x_{n-l(\gamma)}, y_1, \dots, y_{l(\gamma)}) &= \\ &\sum_{0 \leq i \leq n-l(\gamma)} (-1)^i [x_i, f(x_0, \dots, \widehat{x}_i \dots, x_{n-l(\gamma)}, y_1, \dots, y_{l(\gamma)})] \\ &+ \sum_{i < j \leq n-l(\gamma)} (-1)^{j-i-1} f(x_0, \dots, x_{i-1}, [x_i, x_j], x_{i+1}, \dots, \widehat{x}_j \dots, x_{n-l(\gamma)}, y_1, \dots, y_{l(\gamma)}) \\ &+ \sum_{i \leq n-l(\gamma), j} (-1)^{n-l(\gamma)-i+j-1} f(x_0, \dots, \widehat{x}_i \dots, x_{n-l(\gamma)}, y_1, \dots, [x_i, y_j], \dots, y_{l(\gamma)}) \end{aligned}$$

where the x -variables belong to $\mathfrak{g}^\gamma$ and the y 's belong to N^γ . Notice that, as the bracket of $\mathfrak{g}$ is Γ -invariant, N^γ is a $\mathfrak{g}^\gamma$ -submodule of $\mathfrak{g}$. This defines the last summand of ∂_γ^π .

We define a map ψ from the Poisson complex $C_\gamma^\bullet$ to the Koszul complex $CK_\gamma^\bullet$. For any f in C_γ^n , we put :

$$\psi(f) := \Lambda^n \mathfrak{g} \longrightarrow \Lambda^{n-l(\gamma)} \mathfrak{g}^\gamma \otimes \wedge^{l(\gamma)} N^\gamma \xrightarrow{f} S\mathfrak{g}^\gamma \xrightarrow{Sym} \mathcal{U}(\mathfrak{g}^\gamma) \longrightarrow \mathcal{U}(\mathfrak{g})$$

where the first and last map are the usual projection and injection, and Sym is the classical symmetrization map from $S\mathfrak{g}^\gamma$ to $\mathcal{U}(\mathfrak{g}^\gamma)$.

Lemma 4.6. *The map ψ^* is a morphism of complexes.*

Proof. We have to check that ψ commutes with the differentials. For this purpose, let us decompose ∂_γ into a sum of three terms, $\partial_\gamma^1 + \partial_\gamma^2 + \partial_\gamma^3$ corresponding to the three components of its definition. We also use the following decomposition of $\Lambda^{n+1} \mathfrak{g}$ coming from the direct sum $\mathfrak{g} = \mathfrak{g}^\gamma \oplus N^\gamma$:

$$\Lambda^{n+1} \mathfrak{g} = \bigoplus_{p=0}^{l(\gamma)} \Lambda^{n+1-p} \mathfrak{g}^\gamma \otimes \Lambda^p N^\gamma$$

First, as $\psi(f)$ needs $l(\gamma)$ independent variables in N^γ , it follows easily that $\partial_\gamma^1(\psi(f)) = 0$. For the same reason, we check that $\partial(\psi(f))$ is null on the $\Lambda^{n+1-p} \mathfrak{g}^\gamma \otimes \Lambda^p N^\gamma$ whenever $p < l(\gamma)$. Therefore, $\partial(\psi(f))$ reduces to a map from $\Lambda^{n+1-l(\gamma)} \mathfrak{g}^\gamma \otimes \wedge^{l(\gamma)} N^\gamma$ to $\mathcal{U}(\mathfrak{g}^\gamma)$.

To complete the proof, we check that the two formulas of the differentials agree on $\Lambda^{n+1-l(\gamma)}\mathfrak{g}^\gamma \otimes \Lambda^{l(\gamma)}N^\gamma$ thanks to the fact that the symmetrization map Sym is a $\mathfrak{g}$ -morphism. $\square$

Theorem 4.7. *The map ψ is a quasi-isomorphism. Therefore the Hochschild cohomology of $\mathcal{U}(\mathfrak{g}) \rtimes \Gamma$ is isomorphic, as a graded vector space, to the Poisson cohomology of $S(\mathfrak{g}) \rtimes \Gamma$ with the Poisson bracket induced by the Lie bracket of $\mathfrak{g}$:*

$$H^\bullet(\mathcal{U}(\mathfrak{g}) \rtimes \Gamma, \mathcal{U}(\mathfrak{g}) \rtimes \Gamma) \simeq H_\pi^\bullet(S(\mathfrak{g}) \rtimes \Gamma).$$

Proof. Let us introduce a formal parameter $\hbar$ and work over the formal power series $\mathbb{R}[[\hbar]]$. Introduce the $\mathbb{R}[[\hbar]]$ -Lie algebra $\mathfrak{g}_\hbar$ whose underlying $\mathbb{R}[[\hbar]]$ -module is $\mathfrak{g}[[\hbar]]$ with the Lie bracket given by :

$$[x, y]_\hbar = \hbar[x, y],$$

for x and y in $\mathfrak{g}$. Then, the Koszul complex $CK^\bullet(\mathcal{U}(\mathfrak{g}_\hbar) \rtimes \Gamma)$ specializes to that of $S(\mathfrak{g}) \rtimes \Gamma$ for $\hbar = 0$, and, for $\hbar = 1$ to that of $\mathcal{U}(\mathfrak{g}) \rtimes \Gamma$.

Similarly, the Poisson complex $C_\pi^\bullet(\mathcal{U}(\mathfrak{g}_\hbar) \rtimes \Gamma)$ has a zero differential for $\hbar = 0$ and specializes to $C_\pi^\bullet(\mathcal{U}(\mathfrak{g}) \rtimes \Gamma)$ for $\hbar = 1$.

The map ψ extends to the $\mathbb{R}[[\hbar]]$ -context, and defines a morphism of $\mathbb{R}[[\hbar]]$ -complexes. It follows from Subsection 2.1 that ψ specializes to a quasi-isomorphism for $\hbar = 0$. The result is then a consequence of the following standard lemma, which can be found in [8]. $\square$

Lemma 4.8. *Let $C_1^\bullet[[\hbar]]$ and $C_2^\bullet[[\hbar]]$ be two topologically free $\mathbb{R}[[\hbar]]$ -complexes, and ψ a morphism of $\mathbb{R}[[\hbar]]$ -complexes. Suppose that ψ specializes to a quasi-isomorphism for $\hbar = 0$. Then ψ is a quasi-isomorphism.*

We remark that in this paper we have always worked with real coefficient $\mathbb{R}$. Theorem 4.7 is true for a general field $\mathbb{K}$ with characteristic 0.

4.3. Examples of linear Poisson structures. In this section, we provide a large class of linear Poisson structures coming from invariant Lie algebra structures.

We assume that $\mathfrak{g}$ be a Lie algebra and Γ be a finite group acting on $\mathfrak{g}$ preserving its Lie bracket. We choose a Γ invariant metric on $\mathfrak{g}$. Let V be the dual of $\mathfrak{g}$ with the linear Poisson structure π from the Lie bracket. Accordingly, Γ acts on V preserving the Poisson structure π .

For any $\gamma \in \Gamma$ with $l(\gamma) = 2$, let N^γ be the subspace of V normal to V^γ . As $V = V^\gamma \oplus N^\gamma$ and $V^* = V^{\gamma*} \oplus N^{\gamma*}$. One can decompose $\pi = V^* \otimes \wedge^2 V = (V^{\gamma*} \oplus N^{\gamma*}) \otimes (N^\gamma \wedge N^\gamma \oplus N^\gamma \otimes V^\gamma \oplus V^\gamma \otimes N^\gamma \oplus V^\gamma \wedge V^\gamma)$. We define π_γ to be the projection of π onto the component $(V^\gamma)^* \otimes \wedge^2 N^\gamma$.

Proposition 4.9. *The collection $\Pi = \pi + \sum_{\gamma \in l(\gamma)=2} c_\gamma \pi_\gamma$ with constant c_γ satisfying $c_{\alpha\gamma\alpha^{-1}} = c_\gamma$ for any $\alpha \in \Gamma$ defines a linear Poisson structure on $S(V^*) \rtimes \Gamma$.*

Proof. As is explained in subsection 2.1, there is no $Z(\gamma)$ invariant section in $S(V^{\gamma*}) \otimes \wedge^\bullet V^\gamma \otimes \wedge^{l(\gamma)} N^\gamma$ with odd $l(\gamma)$. Furthermore, by Equation (3), any tri-vector field on a γ -component with $l(\gamma) = 4$ is a trivial cocycle. This implies that

$$[\pi_\alpha, \pi_\beta] = 0,$$

if neither α nor β is the identity of Γ .

To prove that Π is a Poisson structure, it is sufficient to prove

$$[\pi, \pi_\gamma] = 0, \quad \gamma \in \Gamma.$$

When γ is identity, the above equation is from the fact that π is the Lie Poisson structure.

When $\gamma \neq 0$, we can decompose π according to the decomposition $V = V^\gamma \oplus N^\gamma$, $V^* = V^{\gamma*} \oplus N^{\gamma*}$. As $\pi \in V^* \otimes V \wedge V$ is γ invariant, π is a sum of the following terms

$$\begin{aligned} \pi_{111} &\in V^{\gamma*} \otimes V^\gamma \wedge V^\gamma, & \pi_{122} &\in V^{\gamma*} \otimes N^\gamma \wedge N^\gamma, \\ \pi_{221} &\in N^{\gamma*} \otimes N^\gamma \wedge V^\gamma, & \pi_{222} &\in N^{\gamma*} \otimes N^\gamma \wedge N^\gamma. \end{aligned}$$

We compute

$$\begin{aligned}
[\pi_{111}, \pi_{111}] &\in V^{\gamma*} \otimes V^{\gamma} \wedge V^{\gamma} \wedge V^{\gamma}, & [\pi_{111}, \pi_{122}] &\in V^{\gamma*} \otimes V^{\gamma} \wedge N^{\gamma} \wedge N^{\gamma}, \\
[\pi_{111}, \pi_{221}] &\in N^{\gamma*} \otimes N^{\gamma} \wedge V^{\gamma} \wedge V^{\gamma}, & [\pi_{111}, \pi_{222}] &= 0, \\
[\pi_{122}, \pi_{122}] &= 0, & [\pi_{122}, \pi_{222}] &\in V^{\gamma*} \otimes N^{\gamma} \wedge N^{\gamma} \wedge N^{\gamma}, \\
[\pi_{122}, \pi_{221}] &\in (V^{\gamma})^* \otimes N^{\gamma} \wedge N^{\gamma} \wedge V^{\gamma} \oplus N^{\gamma*} \otimes N^{\gamma} \wedge N^{\gamma} \wedge N^{\gamma}, \\
[\pi_{221}, \pi_{221}] &\in N^{\gamma*} \otimes N^{\gamma} \wedge V^{\gamma} \wedge V^{\gamma}, & [\pi_{221}, \pi_{222}] &\in N^{\gamma*} \otimes N^{\gamma} \wedge N^{\gamma} \wedge V^{\gamma}, \\
[\pi_{222}, \pi_{222}] &\in N^{\gamma*} \otimes N^{\gamma} \wedge N^{\gamma} \wedge N^{\gamma}.
\end{aligned}$$

According to the fact that $[\pi, \pi] = 0$ and $N^{\gamma} \wedge N^{\gamma} \wedge N^{\gamma} = 0$ for $l(\gamma) = 2$, we see

$$\begin{aligned}
[\pi_{111}, \pi_{111}] &= 0, \\
[\pi_{111}, \pi_{221}] + [\pi_{221}, \pi_{221}] &= 0, \\
[\pi_{111} + \pi_{221}, \pi_{122}] &= 0, \\
[\pi_{122}, \pi_{222}] &= 0, \\
[\pi_{122}, \pi_{122}] &= 0, \\
[\pi_{221}, \pi_{221}] &= 0.
\end{aligned}$$

In particular, this implies $[\pi, \pi_{122}] = 0$. Noticing that $\pi_{\gamma} = \pi_{122}$, we conclude that the Schouten bracket of $[\pi, \pi_{\gamma}] = 0$. $\square$

In the following, we construct explicit examples of linear Poisson structures using Proposition 4.9 on $\mathbb{R}^3$ with the $\mathbb{Z}/2\mathbb{Z} = \{1, e\}$ action by $e(x, y, z) = (-x, -y, z)$. In this case, the fixed point subspace of e is $\{x = y = 0\}$, which is of codimension 2.

Example 4.3.1. Denote $\pi_1 = z\partial_x \wedge \partial_y$. One can easily check that $[\pi_1, \pi_1] = 0$. And the Poisson structure constructed from Proposition 4.9 is $\Pi_1 = \pi_1 + \pi_1|_{x=y=0}e = z\partial_x \wedge \partial_y + z\partial_x \wedge \partial_y|_{x=y=0}e$.

Example 4.3.2. Denote $\pi_2 = z\partial_x \wedge \partial_y + x\partial_x \wedge \partial_z - y\partial_y \wedge \partial_z$. Noticing that $[\partial_x \wedge \partial_y, x\partial_x - y\partial_y] = 0$, we have $[\pi_2, \pi_2] = 0$. And the Poisson structure constructed from Proposition 4.9 is $\Pi_2 = \pi_2 + \pi_2|_{x=y=0}e = z\partial_x \wedge \partial_y + x\partial_x \wedge \partial_z - y\partial_y \wedge \partial_z + z\partial_x \wedge \partial_y|_e$.

We compute the 0-th Poisson cohomology of Π_1 and Π_2 to distinguish them.

- (1) If $f \in (S\mathbb{R}^3)^{\mathbb{Z}/2\mathbb{Z}}$ is a 0-th Poisson cocycle with respect to Π_1 , i.e. $\Pi_1(f) = 0$, then $[\pi_1, f] = 0$ i.e. $\partial_x(f) = \partial_y(f) = 0$, which is also a sufficient condition. Therefore, $H_{\Pi_1}^0 = \{f \in (S\mathbb{R}^3)^{\mathbb{Z}/2\mathbb{Z}} : \partial_x(f) = \partial_y(f) = 0\}$, which means f is a polynomial depending only on z .
- (2) If $f \in (S\mathbb{R}^3)^{\mathbb{Z}/2\mathbb{Z}}$ is a 0-th Poisson cocycle with respect to Π_2 , i.e. $\Pi_2(f) = 0$, then $\pi_2(f) = 0$ and $\pi_1(f)|_{x=y=0} = 0$. This is equivalent to $zf_x + yf_z = 0, zf_y + xf_z = 0, xf_x - yf_y = 0$, and $f_x|_{x=y=0} = f_y|_{x=y=0} = 0$. If we write $f = \sum_{k,m,n} c_{kmn}x^k y^m z^n$, we have that

$$\begin{aligned}
(k+1)c_{k+1mn-1} + (n+1)c_{km-1n+1} &= 0 \\
(m+1)c_{km+1n-1} + (n+1)c_{k-1mn+1} &= 0 \\
kc_{kmn} - mc_{kmn} &= 0 \\
c_{10n} = c_{01n} &= 0.
\end{aligned}$$

From the above third equation, $c_{kmn} = 0$ if $k \neq m$. From the above first and second equation, $(k+1)c_{k+1k+1n} + (n+2)c_{kkn+2} = 0$. From this we can quickly conclude that f is a polynomial of $xy - 1/2z^2$, and $H_{\Pi_2}^0$ consists of polynomials on $xy - 1/2z^2$.

We notice that in the above two examples the 0-th Poisson cohomology of Π_1 (and Π_2) is isomorphic to the 0-th Poisson cohomology of the restriction π_1 (and π_2) of Π_1 (and Π_2) to the identity component of $\mathbb{Z}_2$. In the following proposition, we prove that this is a general phenomena.

Proposition 4.10. *If the group Γ is abelian, the Poisson cohomology of Π is isomorphic to the Poisson cohomology of π .*

Proof. We construct a quasi-isomorphism

$$\Psi : \left(\left(\bigoplus_{\gamma \in \Gamma} S(V^{\gamma*}) \otimes \wedge^{\bullet-l(\gamma)} V^{\gamma} \otimes \wedge^{l(\gamma)} N^{\gamma} \right)^{\Gamma}, [\pi,] \right) \longrightarrow \left(\left(\bigoplus_{\gamma \in \Gamma} S(V^{\gamma*}) \otimes \wedge^{\bullet-l(\gamma)} V^{\gamma} \otimes \wedge^{l(\gamma)} N^{\gamma} \right)^{\Gamma}, [\Pi,] \right).$$

Given $X = \sum_{\gamma} X_{\gamma} \in \left(\bigoplus_{\gamma \in \Gamma} S(V^{\gamma*}) \otimes \wedge^{\bullet-l(\gamma)} V^{\gamma} \otimes \wedge^{l(\gamma)} N^{\gamma} \right)^{\Gamma}$, we define $\Psi(X)$ as a sum of $\sum_{\gamma} \Psi(X_{\gamma})$. Define $\Psi(X_{\gamma})$ as a sum of $1/k! \psi_{\gamma, \alpha_1, \dots, \alpha_k}(X_{\gamma})$ for $k = 0, 1, 2, \dots$. If $l(\gamma \alpha_1 \dots \alpha_k) \neq l(\gamma) + 2k$, then define $\psi_{\gamma, \alpha_1, \dots, \alpha_k}(X_{\gamma}) = 0$; if $l(\gamma \alpha_1 \dots \alpha_k) = l(\gamma) + 2k$, then we define $\psi_{\gamma, \alpha_1, \dots, \alpha_k}(X_{\gamma})$ to be the projection of X_{γ} down to the component

$$\left(S(V^{\gamma \alpha_1 \dots \alpha_k*}) \otimes \wedge^{\bullet} V^{\gamma \alpha_1 \dots \alpha_k} \otimes \wedge^{l(\gamma)} N^{\gamma} \otimes \wedge^2 N^{\alpha_1} \otimes \dots \otimes \wedge^2 N^{\alpha_k} \right)^{\Gamma}.$$

We notice that since $l(\gamma \alpha_1 \dots \alpha_k) \leq \dim(V)$, there are only a finite number of occasions that $\psi_{\gamma, \alpha_1, \dots, \alpha_k}$ is not zero. Therefore, the map $\Psi(X_{\gamma})$ is well-defined.

To prove that Ψ is a chain map, we prove the following equation for $\psi_{\gamma, \alpha_1, \dots, \alpha_k}$ in $H^{\bullet}(S(V^*) \rtimes \Gamma, S(V^*) \rtimes \Gamma)$ at component $\gamma \alpha_1 \dots \alpha_k$,

$$(15) \quad \sum_{\alpha_1 \dots \alpha_k = \delta} \psi_{\gamma, \alpha_1, \dots, \alpha_k}([\pi, X_{\gamma}]) = \sum_{\alpha_1 \dots \alpha_k = \delta} [\pi_{\alpha_k}, k\psi_{\gamma, \alpha_1, \dots, \alpha_{k-1}}(X_{\gamma})] + [\pi, \psi_{\gamma, \alpha_1, \dots, \alpha_{k-1}, \alpha_k}(X_{\gamma})].$$

Equation (15) can be proved by induction. When $k = 0$, the identity is trivial. Assume that Equation (15) is true for $k = n$. We look at the case when $k = n + 1$. Using the induction assumption, we have

$$\begin{aligned} \sum_{\alpha_1 \dots \alpha_{n+1} = \delta} \psi_{\gamma, \alpha_1, \dots, \alpha_{n+1}}([\pi, X_{\gamma}]) &= \sum_{\alpha_1 \dots \alpha_{n+1} = \delta} \psi_{\gamma, \alpha_1, \dots, \alpha_{n+1}}(\psi_{\gamma, \alpha_1, \dots, \alpha_n}([\pi, X_{\gamma}])) \\ &= \sum_{\alpha_1 \dots \alpha_{n+1} = \delta} \psi_{\gamma, \alpha_1, \dots, \alpha_{n+1}} \left(n[\pi_{\alpha_n}, \psi_{\gamma, \alpha_1, \dots, \alpha_{n-1}}(X_{\gamma})] + [\pi, \psi_{\gamma, \alpha_1, \dots, \alpha_n}(X_{\gamma})] \right). \end{aligned}$$

Looking at the contribution of $[\pi, \psi_{\gamma, \alpha_1, \dots, \alpha_n}(X_{\gamma})]$ at $\gamma \alpha_1 \dots \alpha_{n+1}$, we decompose $\psi_{\gamma, \alpha_1, \dots, \alpha_n}(X_{\gamma})$ and π according to $V = V^{\alpha_{n+1}} \oplus N^{\alpha_{n+1}}$. As both π and $\psi_{\gamma, \alpha_1, \dots, \alpha_n}(X_{\gamma})$ are α_{n+1} invariant, we can write $\pi = \pi_0 + \pi_1 + \pi_2$ and $\psi_{\gamma, \alpha_1, \dots, \alpha_n}(X_{\gamma}) = X_0 + X_1 + X_2$ such that π_i, X_i contain i number of vector fields along $N^{\alpha_{n+1}}$. It is easy to see that $[\pi_0 + \pi_1, X_0 + X_1]$ will contribute zero at $\gamma \alpha_1 \dots \alpha_{n+1}$ by the facts that π is a linear Poisson structure and $[\pi_0 + \pi_1, X_0 + X_1]$ at $\gamma \alpha_1 \dots \alpha_{n+1}$ is the restriction to $V^{\gamma \alpha_1 \dots \alpha_{n+1}}$ of the component

$$S(V^{\gamma \alpha_1 \dots \alpha_{n+1}*}) \otimes \wedge^{\bullet} V^{\gamma \alpha_1 \dots \alpha_{n+1}} \otimes \wedge^{l(\gamma)} N^{\gamma} \otimes \wedge^2 N^{\alpha_1} \otimes \dots \otimes \wedge^2 N^{\alpha_{n+1}}$$

in $[\pi_0 + \pi_1, X_0 + X_1]$. Furthermore, $[\pi_2, X_2]$ vanishes as it is a 3 vector field normal to $V^{\alpha_{n+1}}$, which is of codimension 2. Therefore, $[\pi, \psi_{\gamma, \alpha_1, \dots, \alpha_n}(X_{\gamma})]$ at $\gamma \alpha_1 \dots \alpha_{n+1}$ is equal to

$$[\pi, X_2] + [\pi_2, X] = [\pi, \psi_{\gamma, \alpha_1, \dots, \alpha_{n+1}}(X_{\gamma})] + [\pi_{\alpha_{n+1}}, \psi_{\gamma, \alpha_1, \dots, \alpha_n}(X_{\gamma})].$$

Next, we observe that $\psi_{\gamma, \alpha_1, \dots, \alpha_{n+1}}([\pi_{\alpha_n}, \psi_{\gamma, \alpha_1, \dots, \alpha_{n-1}}(X_{\gamma})])$ is equal to $[\pi_{\alpha_n}, \psi_{\gamma, \alpha_1, \dots, \alpha_{n-1}, \alpha_{n+1}}(X_{\gamma})]$ as the space N^{α_n} and $N^{\alpha_{n+1}}$ are orthogonal to each other by the assumption that $l(\gamma) + 2n + 2 = l(\gamma \alpha_1 \dots \alpha_{n+1})$. Hence, we have the above Equation (15) for $k = n + 1$.

Using Equation (15), we can easily check that Ψ is a chain map. To prove that Ψ is a quasi-isomorphism, we look at the filtration with respect to the grading $l(\gamma)$ as is used in the proof of Proposition 4.2. It is straight forward to see that the induced chain map of Ψ is identity at E_1 as for $k = 0$, $\psi_{\gamma}(X_{\gamma}) = X_{\gamma}$, which implies that Ψ is a quasi-isomorphism. $\square$

5. EXAMPLES OF QUANTIZATION

In this section, we study some examples of quantization of constant Poisson structures. We look at $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ action on $\mathbb{R}^2$ by the rotation

$$\gamma : (x, y) \mapsto \left(\cos\left(\frac{2\pi}{n}\right)x - \sin\left(\frac{2\pi}{n}\right)y, \sin\left(\frac{2\pi}{n}\right)x + \cos\left(\frac{2\pi}{n}\right)y \right), \quad \gamma^n = 1,$$

where γ is the generator of $\mathbb{Z}/n\mathbb{Z}$. If we introduce complex coordinates $z = x + iy, \bar{z} = x - iy$, then the above action is diagonalized

$$\gamma : (z, \bar{z}) \mapsto (\exp(\frac{2\pi i}{n})z, \exp(-\frac{2\pi i}{n})\bar{z}).$$

We study the Poisson structure of the following form $\pi : \wedge^2 \mathbb{R}^2 \rightarrow \mathbb{R}\mathbb{Z}_n$ by $\pi(x, y) = \gamma$. In complex coordinates $\pi(z, \bar{z}) = -i/2\gamma$. By [9][Corollary 4.2], π defines a noncommutative structure on $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$, and by Proposition 4.2, π can be quantized. In the following two sections, we study properties of quantization of π .

5.1. A Moyal type formula. In this subsection, we provide an explicit formula for quantization of π , which is a generalization of Moyal product. We would like to point out that many of the following formulas appear already in Nadaud's paper [12]. We prove that in the case of finite group, this product is convergent. We start with introducing several operators on $S(\mathbb{R}^2)$. We work with complex coordinates $z = x + iy, \bar{z} = x - iy$.

Define $D_z, D_{\bar{z}} : S(\mathbb{R}^2) \rightarrow S(\mathbb{R}^2)$ as

$$D_z(f) = \frac{f(z, \bar{z}) - f(e^{\frac{2\pi i}{n}}z, \bar{z})}{(1 - e^{\frac{2\pi i}{n}})z}, \quad D_{\bar{z}}(f) = \frac{f(z, \bar{z}) - f(z, e^{-\frac{2\pi i}{n}}\bar{z})}{(1 - e^{-\frac{2\pi i}{n}})\bar{z}}.$$

Define $\sigma_z, \sigma_{\bar{z}} : S(\mathbb{R}^2) \rightarrow S(\mathbb{R}^2)$ as

$$\sigma_z(f)(z, \bar{z}) = f(e^{\frac{2\pi i}{n}}z, \bar{z}) \quad \sigma_{\bar{z}}(f)(z, \bar{z}) = f(z, e^{-\frac{2\pi i}{n}}\bar{z}).$$

Let $q = \exp(2\pi i/n)$. Define $[k]_q = 1 + q + \dots + q^{k-1}$. Define the following star product $\star$ on $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$. Define $f\gamma^k \star g\gamma^l$ by

$$f\gamma^k \star g\gamma^l = \sum_{j=0}^{\infty} \frac{(\frac{i\hbar}{2})^j}{[j]_q!} (D_z)^j(f)(\sigma_z D_z)^j(\gamma^k(g))\gamma^{j+k+l}$$

To prove the associativity of $\star$, we study properties of $D_z, D_{\bar{z}}$.

Lemme 5.1.

$$D_z^k(fg) = \sum_{i=0}^k \frac{[k]_q!}{[k-i]_q![i]_q!} D_z^i(f)\sigma_z^i D_z^{k-i}(g), \quad k \geq 0.$$

Proof. We prove this by induction. When $k = 0$, this identity is trivial.

Assume that this identity holds for k . For $k+1$, we compute

$$\begin{aligned} D_z^{k+1}(fg) &= D_z \left(\sum_{i=0}^k \frac{[k]_q!}{[k-i]_q![i]_q!} D_z^i(f)\sigma_z^i D_z^{k-i}(g) \right) \\ &= \sum_{i=0}^k \frac{[k]_q!}{[k-i]_q![i]_q!} D_z(D_z^i(f)\sigma_z^i D_z^{k-i}(g)) \\ &= \sum_{i=0}^k \left(\frac{[k]_q!}{[k-i]_q![i]_q!} D_z^{i+1}(f)\sigma_z^{i+1} D_z^{k-i}(g) + \frac{[k]_q!}{[k-i]_q![i]_q!} D_z^i(f)D_z\sigma_z^i D_z^{k-i}(g) \right) \\ &= D_z^{k+1}(f)\sigma_z^{k+1}(g) + fD_z^{k+1}(g) \\ &+ \sum_{i=1}^k \left(\frac{[k]_q!}{[k-i+1]_q![i-1]_q!} D_z^i(f)\sigma_z^i D_z^{k-i+1}(g) + \frac{[k]_q!}{[k-i]_q![i]_q!} q^i D_z^i(f)\sigma_z^i D_z^{k-i+1}(g) \right) \\ &= D_z^{k+1}(f)\sigma_z^{k+1}(g) + \sum_{i=1}^k \frac{[k+1]_q!}{[k-i+1]_q![i]_q!} D_z^i(f)\sigma_z^i D_z^{k-i+1}(g) + fD_z^{k+1}(g) \\ &= \sum_{i=0}^{k+1} \frac{[k+1]_q!}{[k-i+1]_q![i]_q!} D_z^i(f)\sigma_z^i D_z^{k+1-i}(g). \end{aligned}$$

□

We start to prove the associativity of $\star$.

$$\begin{aligned}
& (f \star g) \star h \\
&= \sum_k \left(\frac{i\hbar}{2}\right)^k \frac{1}{[k]_q!} D_z^k(f) \sigma_z^k D_{\bar{z}}^k(g) \gamma^k \star h \\
&= \sum_{k,l} \left(\frac{i\hbar}{2}\right)^{k+l} \frac{1}{[l]_q! [k]_q!} D_z^l \left(D_z^k(f) \sigma_z^k D_{\bar{z}}^k(g) \right) \sigma_z^l D_{\bar{z}}^l(\gamma^k(h)) \gamma^{k+l}
\end{aligned}$$

Using Lemma 5.1, we have that the above product is equal to

$$\begin{aligned}
&= \sum_{k,l} \left(\frac{i\hbar}{2}\right)^{k+l} \frac{1}{[l]_q! [k]_q!} \frac{[l]_q!}{[l-m]_q! [m]_q!} D_z^{m+k}(f) \sigma_z^m D_z^{l-m} \sigma_z^k D_{\bar{z}}^k(g) \sigma_z^l D_{\bar{z}}^l(\gamma^k(h)) \gamma^{k+l} \\
&= \sum_{s,t,k} \left(\frac{i\hbar}{2}\right)^{s+t+l} \frac{1}{[k]_q! [s]_q! [t]_q!} D_z^{s+k}(f) q^{kt} \sigma_z^{s+k} D_z^t D_{\bar{z}}^k(g) \sigma_z^{s+t} D_{\bar{z}}^{s+t}(\gamma^k(h)) \gamma^{k+s+t} \\
&= \sum_{s,t,k} \left(\frac{i\hbar}{2}\right)^{s+t+l} \frac{1}{[k]_q! [s]_q! [t]_q!} D_z^{s+k}(f) q^{kt} \sigma_z^{s+k} D_z^t D_{\bar{z}}^k(g) \sigma_z^{s+t} D_{\bar{z}}^{s+t} \sigma_z^k \sigma_{\bar{z}}^k(h) \gamma^{k+s+t} \\
&= \sum_{s,t,k} \left(\frac{i\hbar}{2}\right)^{s+t+l} \frac{1}{[k]_q! [s]_q! [t]_q!} D_z^{s+k}(f) q^{kt} \sigma_z^{s+k} D_z^t D_{\bar{z}}^k(g) \sigma_z^{s+t+k} \sigma_{\bar{z}}^k q^{-k(s+t)} D_{\bar{z}}^{s+t}(h) \gamma^{k+s+t}
\end{aligned}$$

On the other hand, we compute

$$\begin{aligned}
& f \star (g \star h) \\
&= \sum_k \left(\frac{i\hbar}{2}\right)^k \frac{1}{[k]_q!} f \star (D_z^k(g) \sigma_z^k D_{\bar{z}}^k(h) \gamma^k) \\
&= \sum_{k,l} \left(\frac{i\hbar}{2}\right)^{k+l} \frac{1}{[l]_q! [k]_q!} D_z^l(f) \sigma_z^l D_{\bar{z}}^l \left(D_z^k(g) \sigma_z^k D_{\bar{z}}^k(h) \right) \gamma^{k+l}
\end{aligned}$$

Applying the similar formula for D_z as Lemma 5.1, we have

$$\begin{aligned}
&= \sum_{k,s,t} \left(\frac{i\hbar}{2}\right)^{k+s+t} \frac{1}{[k]_q! [s+t]_q!} \frac{[s+t]_{q^{-1}}!}{[s]_{q^{-1}}! [t]_{q^{-1}}!} D_z^{s+t}(f) \sigma_z^{s+t} D_{\bar{z}}^s D_z^k(g) \sigma_z^{s+t} \sigma_{\bar{z}}^s D_{\bar{z}}^t \sigma_z^k D_{\bar{z}}^k(h) \gamma^{k+s+t} \\
&= \sum_{k,s,t} \left(\frac{i\hbar}{2}\right)^{k+s+t} \frac{1}{[k]_q! [s+t]_q!} \frac{[s+t]_{q^{-1}}!}{[s]_{q^{-1}}! [t]_{q^{-1}}!} D_z^{s+t}(f) \sigma_z^{s+t} D_{\bar{z}}^s D_z^k(g) \sigma_z^{s+t+k} \sigma_{\bar{z}}^s D_{\bar{z}}^{t+k}(h) \gamma^{k+s+t}
\end{aligned}$$

In the above equation, we make the change $s \mapsto k, t \mapsto s, k \mapsto t$, then we have

$$\sum_{k,s,t} \left(\frac{i\hbar}{2}\right)^{k+s+t} \frac{1}{[t]_q! [k+s]_q!} \frac{[k+s]_{q^{-1}}!}{[k]_{q^{-1}}! [s]_{q^{-1}}!} D_z^{k+s}(f) \sigma_z^{k+s} D_{\bar{z}}^k D_z^t(g) \sigma_z^{s+t+k} \sigma_{\bar{z}}^k D_{\bar{z}}^{s+t}(h) \gamma^{k+s+t}$$

which is identified with the above expression of $(f \star g) \star h$. We remark that $[k]_q = [k]_{q^{-1}} q^{k-1}$ and $[k]_q! = [k]_{q^{-1}}! q^{(k-1)k/2}$. Therefore, we conclude that $\star$ defines an associative deformation of $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$, whose $\hbar$ component is equal to $i/2 D_z \otimes \sigma_z D_{\bar{z}}$, which is cohomologous to the Poisson structure π .

We remark that our proof of associativity of $\star$ is slightly different from [12]. One can view $f \star g$ as an extension of the Moyal product as follows

$$f \star g = m \circ \exp_q \left(\frac{i\hbar}{2} (D_z \otimes \sigma_z D_{\bar{z}} \otimes \gamma) \right) (f \otimes g \otimes 1).$$

Nadaud proved the associativity of $\star$ analogous to the associativity of Moyal product by using the property of the q -exponential. Here our proof is more straightforward, and it leads to the following more precise formula for $\star$.

We have the following property for the operator D_z .

Proposition 5.2.

$$D_z^m(f) = \frac{\sum_{i=0}^m (-1)^{m-i} \frac{[m]_q!}{[m-i]_q! [i]_q!} q^{i(i-1)/2} \sigma_z^{m-i} f}{(1-q)^m q^{m(m-1)/2} z^m}.$$

In particular, when $m = n$, $D_z^n(f) = 0$ for any f . And this implies that

$$f \star g = \sum_{j=0}^{n-1} \frac{(\frac{i\hbar}{2})^j}{[j]_q!} D_z^j(f) \sigma_z^j D_z^j(g) \gamma^j.$$

Proof. We prove the identity by induction. When $m = 1$, we have

$$D_z(f) = \frac{f(z, \bar{z}) - f(qz, \bar{z})}{z - \gamma(z)} = \frac{f(z, \bar{z}) - \sigma_z(f)(z, \bar{z})}{(1-q)z}.$$

Assume that the above identity holds for m . Then for $m+1$,

$$\begin{aligned} D_z^{m+1}(f) &= D_z(D_z^m(f)) = D_z\left(\frac{\sum_{i=0}^m (-1)^{m-i} \frac{[m]_q!}{[m-i]_q! [i]_q!} q^{i(i-1)/2} \sigma_z^{m-i} f}{(1-q)^m q^{m(m-1)/2} z^m}\right) \\ &= \sum_{i=0}^m (-1)^{m-i} \frac{[m]_q!}{[m-i]_q! [i]_q!} \frac{q^{i(i-1)/2}}{(1-q)^m q^{m(m-1)/2}} D_z\left(\frac{\sigma_z^{m-i} f}{z^m}\right) \\ &= \sum_{i=0}^m (-1)^{m-i} \frac{[m]_q!}{[m-i]_q! [i]_q!} \frac{q^{i(i-1)/2}}{(1-q)^m q^{m(m-1)/2}} \frac{\frac{\sigma_z^{m-i} f}{z^m} - \frac{\sigma_z^{m-i+1}(f)}{\gamma(z^m)}}{(z - \gamma(z))} \\ &= \sum_{i=0}^m (-1)^{m-i} \frac{[m]_q!}{[m-i]_q! [i]_q!} \frac{q^{i(i-1)/2}}{(1-q)^m q^{m(m-1)/2}} \frac{q^m \sigma_z^{m-i}(f) - \sigma_z^{m-i+1}(f)}{q^m (1-q) z^{m+1}} \\ &= \frac{1}{q^{m(m+1)/2} (1-q)^{m+1} z^{m+1}} \sum_{i=0}^m (-1)^i \frac{[m]_q!}{[m-i]_q! [i]_q!} q^{(m-i)(m-i-1)/2} (q^m \sigma_z^i(f) - \sigma_z^{i+1}(f)) \\ &= \frac{1}{q^{m(m+1)/2} (1-q)^{m+1} z^{m+1}} \left(q^{m(m+1)/2} f + (-1)^{m+1} \sigma_z^{m+1}(f) + \right. \\ &\quad \left. + [m]_q! \sum_{i=1}^m (-1)^i \left(\frac{q^m q^{(m-i)(m-i-1)/2}}{[m-i]_q! [i]_q!} + \frac{q^{(m+1-i)(m-i)/2}}{[m-i+1]_q! [i-1]_q!} \right) \sigma_z^i(f) \right) \\ &= \frac{1}{q^{m(m+1)/2} (1-q)^{m+1} z^{m+1}} \left(q^{m(m+1)/2} f + (-1)^{m+1} \sigma_z^{m+1}(f) + \right. \\ &\quad \left. + [m]_q! \sum_{i=1}^m (-1)^i \frac{q^{(m+1-i)(m-i)/2} (q^i [m-i+1]_q! + [i]_q!)}{[m+1-i]_q! [i]_q!} \sigma_z^i(f) \right) \\ &= \frac{1}{q^{m(m+1)/2} (1-q)^{m+1} z^{m+1}} \sum_{i=0}^{m+1} (-1)^i q^{(m+1-i)(m-i)/2} \frac{[m+1]_q!}{[m+1-i]_q! [i]_q!} \sigma_z^i(f). \end{aligned}$$

We have proved the identity of $D_z^m(f)$ by induction. To conclude that $D_z^n(f) = 0$. We see that by the above formula of $D_z^n(f)$, as $[n]_q = 0$,

$$D_z^n(f) = \frac{1}{q^{n(n-1)/2} (1-q)^n z^n} (q^{(n-1)n/2} f + (-1)^n \sigma_z^n(f)).$$

Since $\sigma_z^n(f) = f$, we have

$$\frac{1}{q^{n(n-1)/2} (1-q)^n z^n} (q^{(n-1)n/2} + (-1)^n)(f).$$

The statement follows from the identity $q^{(n-1)n/2} + (-1)^n = 0$. $\square$

We conclude from Proposition 5.2 that the star product $\star$ on $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$ is convergent for any value of $\hbar$.

In particular, when $n = 2$, we have the following explicit formula of a deformation on $S(\mathbb{R}^2) \rtimes \mathbb{Z}_2$

$$f \star g = fh + \left(\frac{i\hbar}{2}\right) \frac{f(z, \bar{z}) - f(-z, \bar{z})}{2z} \frac{f(-z, \bar{z}) - f(-z, -\bar{z})}{2\bar{z}}.$$

Remark 5.1.1. Here our formula of product uses “normal ordering”, by which we mean that D_z is contained only in the left component and $D_{\bar{z}}$ is contained in the right component. We can also define product with “anti-normal ordering” or “symmetric ordering” as [12]. The similar property like Proposition 5.2 extends directly.

Remark 5.1.2. We observe that the formula for the star product $\star$ on $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n[[\hbar]]$ works well for the algebra $C_c^\infty(\mathbb{R}^2) \rtimes \mathbb{Z}_n[[\hbar]]$. Again, $\star$ product is convergent for any two smooth functions f and g on $\mathbb{R}^2$.

5.2. Deformation of singularity. In this subsection, we compute the center of the above quantized algebra $(S(\mathbb{R}^2) \rtimes \mathbb{Z}_n[[\hbar]], \star)$. We prove that the center is closely connected to the deformation of the underlying quotient space $\mathbb{R}^2/\mathbb{Z}_n$. We must say that this kind of idea is already in [6]. Here we are giving concrete examples about this idea.

We write an element in $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n[[\hbar]]$ by $\sum_{i=0}^{n-1} f_i \gamma^i$ with f_i in $S(\mathbb{R}^2)[[\hbar]]$.

Proposition 5.3. If $f = \sum_{i=0}^{n-1} f_i \gamma^i$ is in the center of $(S(\mathbb{R}^2) \rtimes \mathbb{Z}_n[[\hbar]], \star)$, then f is completely determined by f_0 by the following formula

$$\gamma(f_0) = f_0, \quad f_j = \left(\frac{i\hbar}{2}\right)^j \frac{D_z^j(f_0)}{[j]_q!(1-q^{-1})^j \bar{z}^j} = \left(-\frac{i\hbar}{2}\right)^j q^{-j(j-1)/2} \frac{\sigma_z^j D_{\bar{z}}^j(f_0)}{(1-q)^j [j]_q! z^j}, \quad i = 1, \dots, n-1.$$

Therefore, as a vector space the center of the quantum algebra is isomorphic to $S(\mathbb{R}^2)^{\mathbb{Z}_n}$, the algebra of $\mathbb{Z}_n$ invariant polynomials on V .

Proof. We need to first prove the above two expressions for f_i are same. We prove this using Proposition 5.2.

$$\begin{aligned} & \left(-\frac{i\hbar}{2}\right)^j q^{-j(j-1)/2} \frac{\sigma_z^j D_{\bar{z}}^j(f_0)}{(1-q)^j [j]_q! z^j} \\ &= \frac{\left(-\frac{i\hbar}{2}\right)^j q^{-j(j-1)/2}}{(1-q)^j [j]_q! z^j} \sigma_z^j \frac{\sum_{k=0}^j (-1)^k \frac{[j]_q!}{[j-k]_q! [k]_q!} q^{-(j-k)(j-k-1)/2} \sigma_{\bar{z}}^k(f_0)}{(1-q^{-1})^j q^{-j(j-1)/2} \bar{z}^j} \\ &= \frac{\left(\frac{i\hbar}{2}\right)^j q^{-j(j-1)/2} \sum_k (-1)^{j-k} \frac{1}{[j-k]_q! [j]_q!} q^{(j-k)(j-k-1)/2 + k(k-1)/2 - (j-k)(j-k-1)/2} \sigma_z^j \sigma_{\bar{z}}^k(f_0)}{(1-q)^j (1-q^{-1})^j z^j \bar{z}^j} \\ &= \frac{\left(\frac{i\hbar}{2}\right)^j q^{-j(j-1)/2} \sum_k (-1)^{j-k} \frac{[j]_q!}{[j-k]_q! [k]_q!} q^{k(k-1)} \sigma_z^{j-k} f_0}{[j]_q! (1-q^{-1})^j \bar{z}^j} \frac{1}{(1-q)^j z^j} \end{aligned}$$

where in the last line we have used that $\sigma_z \sigma_{\bar{z}}(f_0) = \gamma(f_0) = f_0$. By Proposition 5.2, we conclude that the above expression is equal to

$$\frac{\left(\frac{i\hbar}{2}\right)^j}{[j]_q! (1-q^{-1})^j \bar{z}^j} D_z^j(f_0).$$

Let $f = f_0 + f_1 \gamma + \dots + f_{n-1} \gamma^{n-1}$ be an element in the center of $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n$. We compute $f \star z = \sum_i f_i \star \gamma^i(z) \gamma^i = \sum_i q^i f_i z \gamma^i$, and $z \star f = \sum_j z \star f_j \gamma^j = (\sum_j z f_j + \frac{i\hbar}{2} \sigma_z D_{\bar{z}}(f_j) \gamma) \gamma^j = \sum_j (z f_j + \frac{i\hbar}{2} \sigma_z D_{\bar{z}}(f_{j-1}) \gamma) \gamma^j$. As $f \star z = z \star f$, $f_j = -\frac{i\hbar}{2} \sigma_z D_{\bar{z}}(f_{j-1}) / [(1-q)[j]_q! z]$. And we can solve by induction to find that f_i has to be the form expressed in the Proposition.

To prove that the above defined $f = f_0 + f_1 \gamma + \dots + f_{n-1} \gamma^{n-1}$ is in the center, we show in the following $f \star g = g \star f$ for any $g \in S(\mathbb{R}^2)$ and $\gamma(f) = f$.

For $\gamma(f) = f$, it is enough to prove that $\gamma(f_i) = f_i$. This is obvious from the following identity

$$f_j = \frac{\left(\frac{i\hbar}{2}\right)^j q^{-j(j-1)/2}}{[j]_q! (1-q^{-1})^j \bar{z}^j} \sum_k (-1)^{j-k} \frac{[j]_q!}{[j-k]_q! [k]_q!} q^{k(k-1)} \sigma_z^{j-k} f_0$$

and the fact that f_0 is γ invariant.

For $f \star g = g \star f$, we compute the two sides of the equation separately.

$$\begin{aligned}
f \star g &= \sum_i f_i \gamma^i \star g = \sum_i f_i \star \gamma^i(g) \gamma^i \\
&= \sum_{i,j} \frac{(\frac{i\hbar}{2})^j}{[j]_q!} D_z^j(f_i) \sigma_z^j D_z^j(\gamma^i(g)) \gamma^{i+j} \\
&= \sum_{j,k} \frac{(\frac{i\hbar}{2})^j}{[j]_q!} D_z^j \left(\left(\frac{i\hbar}{2} \right)^k \frac{D_z^k(f_0)}{[k]_q! (1-q^{-1})^k \bar{z}^k} \right) \sigma_z^j D_z^j(\gamma^k(g)) \gamma^{j+k} \\
&= \sum_{j,k} \frac{(\frac{i\hbar}{2})^{j+k}}{[k]_q! [j]_q!} \frac{D_z^{j+k}(f_0) \sigma_z^j D_z^j(\gamma^j(g)) \gamma^{j+k}}{(1-q^{-1})^k \bar{z}^k} \\
&= \sum_k \left(\frac{i\hbar}{2} \right)^k D_z^k(f_0) \sum_{j=0}^k \frac{\sigma_z^{k-j} D_z^{k-j}(\gamma^j(g)) \gamma^k}{[j]_q! [k-j]_q! (1-q^{-1})^j \bar{z}^j}.
\end{aligned}$$

Applying Proposition 5.2 to the above D_z^{k-j} , we have that $f \star g$ is equal to

$$\begin{aligned}
&= \sum_k \left(\frac{i\hbar}{2} \right)^k D_z^k(f_0) \sum_{j=0}^k \frac{1}{[j]_q! [k-j]_q! (1-q^{-1})^j \bar{z}^j} \\
&\quad \sigma_z^{k-j} \left(\frac{\sum_{l=0}^{k-j} (-1)^l \frac{[k-j-l]_{q^{-1}}!}{[k-j-l]_{q^{-1}}! [l]_{q^{-1}}!} q^{-(k-j-l)(k-j-l-1)/2} \sigma_z^l (\sigma_z^j \sigma_z^j g)}{(1-q^{-1})^{k-j} q^{-(k-j)(k-j-1)/2} \bar{z}^{k-j}} \right) \gamma^k \\
&= \sum_k \left(\frac{i\hbar}{2} \right)^k D_z^k(f_0) \sum_{j=0}^k \sum_{l=0}^j \frac{(-1)^{j-l} q^{(j-l)(j-l-1)/2} \sigma_z^k \sigma_z^j g}{[l]_q! [k-j]_q! [j-l]_q! (1-q^{-1})^k \bar{z}^k} \gamma^k.
\end{aligned}$$

It is not difficult to check that $\sum_{i=0}^j \frac{(-1)^{j-i} q^{(j-i)(j-i-1)/2}}{[i]_q! [j]_q!} = 0$ if $j \neq 0$, and $= 1$ if $j = 0$. We replace this computation into the above line and have

$$f \star g = \sum \frac{(\frac{i\hbar}{2})^k}{[k]_q! (1-q^{-1})^k \bar{z}^k} D_z^k(f_0) \sigma_z^k g \gamma^k.$$

The computation of $g \star f$ is similar to the above and we conclude that

$$f \star g = \sum \frac{(\frac{i\hbar}{2})^k}{[k]_q! (1-q^{-1})^k \bar{z}^k} D_z^k(f_0) \sigma_z^k g \gamma^k = g \star f.$$

□

Remark 5.2.1. The above proof on f belonging to the center of quantum algebra can be simplified by checking that f commutes with the generators of $S(\mathbb{R}^2) \rtimes \mathbb{Z}_n[[\hbar]]$, which consists of $z, \bar{z}, \gamma$. We have taken the above proof because it extends to the algebra $C_c^\infty(\mathbb{R}^2) \rtimes \mathbb{Z}_n[[\hbar]]$ directly.

In the following, we study the algebraic structure on the center $Z_{\mathbb{C}}^h(\mathbb{R}^2, \mathbb{Z}_n)$ of the (complexified) quantum algebra $(S(\mathbb{R}^2) \rtimes \mathbb{Z}_n[[\hbar]]) \otimes_{\mathbb{R}} \mathbb{C}$. (The reason that we consider the complexified algebra is that it is relatively easy to write down a set of generators and their relations for the center. However, our following discussion also works for the real algebra.) It is easy to check that $u = z^n$, $v = \bar{z}^n$ and $w = z\bar{z} + \frac{i\hbar}{1-q^{-1}}\gamma$ are in the center of the complexified quantum algebra $(S(\mathbb{R}^2) \rtimes \mathbb{Z}_n[[\hbar]]) \otimes_{\mathbb{R}} \mathbb{C}$. According to the isomorphism as vector space between the center of the quantum algebra and $S(\mathbb{R}^2)^\Gamma$, we know that $u = z^n$, $v = \bar{z}^n$ and $w = z\bar{z} + i\hbar\gamma/(2(1-q^{-1}))$ generates the whole center $Z_{\mathbb{C}}^h(\mathbb{R}^2, \mathbb{Z}_n)$. The relation

between these three generators is generated by

$$\begin{aligned}
z^n \star \bar{z}^n &= z \star \cdots \star z \star \bar{z} \star \cdots \star \bar{z} = z^{\star(n-1)}(z\bar{z} + \frac{i\hbar}{2}\gamma)\bar{z}^{\star(n-1)} \\
&= z^{\star(n-1)}(w + \frac{i\hbar}{1-q}\gamma)\bar{z}^{\star(n-1)} \\
&= z^{\star(n-1)} \star \bar{z}^{\star(n-1)} \star (w + \frac{i\hbar}{1-q}\gamma) \\
&= z^{\star(n-2)} \star \bar{z}^{\star(n-2)} \star (w + \frac{i\hbar}{1-q}\gamma) \star (w + \frac{i\hbar}{1-q}\gamma) \\
&\quad \dots \\
&= (w + \frac{i\hbar}{1-q}\gamma) \star \cdots \star (w + \frac{i\hbar}{1-q}\gamma)
\end{aligned}$$

As w is in the center, the last line can be viewed as the expansion of $w^{\star(n)} + \frac{(\frac{i\hbar}{2})^n q^{-\frac{n(n-1)}{2}}}{(1-q)^n} \gamma^n$. Therefore, $Z_{\mathbb{C}}^{\hbar}(\mathbb{R}^2, \mathbb{Z}_n)$ is generated by u, v, w with the relation that $u^n v^n = w^n + (\frac{i\hbar}{2})^n q^{-\frac{n(n-1)}{2}} / (1-q)^n$. In particular, we define a deformation of the cone $\langle u, v, w \rangle / \{u^n v^n = w^n\}$, which is the algebra of polynomials on the quotient $V/\mathbb{Z}_n$. Furthermore, we notice that for function $F^{\hbar}(u, v, w) = u^n v^n - w^n - \frac{(\frac{i\hbar}{2})^n q^{-\frac{n(n-1)}{2}}}{(1-q)^n}$, $(F_u^{\hbar}, F_v^{\hbar}, F_w^{\hbar})$ is a non-zero vector in $\mathbb{C}^3$ if and only if $u = v = w = 0$, which is not on the surface determined by $F^{\hbar} = 0$. Therefore, we conclude that $F^{\hbar} = 0$ determines a smooth surface when $\hbar \neq 0$, and $Z_{\mathbb{C}}^{\hbar}(\mathbb{R}^2, \mathbb{Z}_n)$ is a nontrivial deformation of the cone $\mathbb{C}^2/\mathbb{Z}_n$.

On the other hand, we look at the 0-th Poisson cohomology of the Poisson structure π_{γ} on $H^{\bullet}(S(\mathbb{R}^2) \rtimes \mathbb{Z}_n, S(\mathbb{R}^2) \rtimes \mathbb{Z}_n) \otimes_{\mathbb{R}} \mathbb{C}$. It is not difficult to see that the 0-th Poisson cohomology $H_{\pi_{\gamma}}^0$ is isomorphic to $S(\mathbb{R}^2)^{\mathbb{Z}_n} \otimes_{\mathbb{R}} \mathbb{C}$ as an algebra.

We summarize the above study into the following corollary.

Proposition 5.4. *The center $Z_{\mathbb{C}}^{\hbar}(\mathbb{R}^2, \mathbb{Z}_n)$ is not isomorphic to the Poisson center $H_{\pi_{\gamma}}^0[[\hbar]] = S(\mathbb{R}^2)^{\mathbb{Z}_n}[[\hbar]]$, but defines a nontrivial deformation.*

In particular, when $n = 2$, the center $Z_{\mathbb{C}}^{\hbar}(\mathbb{R}^2, \mathbb{Z}_2)$ is equal to $\langle u, v, w \rangle / \{uv = w^2 + \frac{\hbar^2}{16}\}$. This is the algebra of polynomial functions on the hyperboloid (when $\hbar$ is real, the surface is one-sheeted, when $\hbar$ is imaginary, the surface is two-sheeted.).

Remark 5.2.2. *We can extend the above discussion of center to quantization of more general Poisson structures. For example, the same discussion holds true for the center of the quantization of the linear Poisson structure $z\partial_x \wedge \partial_y$ on $S(\mathbb{R}^3) \times \mathbb{Z}_n$ with $\mathbb{Z}_n$ acting on $\mathbb{R}^3$ by rotating the x, y -plane and fixing the z axis.*

In summary, we have seen that the center of the quantization of a Poisson structure π on an orbifold may not be isomorphic to the 0-th Poisson cohomology of π as an algebra. On the other hand, the well-known Duflo's isomorphism for a Lie algebra states that the center of the universal enveloping algebra $\mathcal{U}(\mathfrak{g})$ is isomorphic to the Poisson center of $S(\mathfrak{g})$ as an algebra. Our examples suggest that the natural extension of Duflo's isomorphism does not hold in the case of quantization of Lie Poisson structures on orbifolds. We plan to study this interesting phenomena in future publications.

REFERENCES

- [1] J. Alev, M. Farinati, T. Lambre and A. Solotar, *Homologie des invariants d'une algèbre de Weyl sous l'action d'un groupe fini*, J. of Algebra, 232, 564-577 (2000).
- [2] J. Block, and E. Getzler, *Quantization of foliations*, Proceedings of the XXth International Conference on Differential Geometric Methods in Theoretical Physics, 1991, New York City, Vol. 1-2, World Scientific (Singapore), 471-487 (1992).
- [3] A. Braverman, D. Gaitsgory, *Poincar-Birkhoff-Witt theorem for quadratic algebras of Koszul type*, J. Algebra **181**, 315-328 (1996).
- [4] J.L. Brylinski, V. Nistor, *Cyclic cohomology of étale groupoids*, K-Theory 8 (1994), no. 4, 341-365.

- [5] L. Dornhoff, *Group representation theory*, Part A: Ordinary representation theory, Pure and Applied Mathematics, 7. Marcel Dekker, Inc., New York, 1971, 1–254.
- [6] P. Etingof, V. Ginzburg, *Symplectic reflection algebras, Calogero-Moser space and deformed Harish-Chandra homomorphism*, Invent. Math. **147**, 243–348 (2002).
- [7] E. Getzler, *Cartan homotopy formulas and Gauss-Manin connection in cyclic homology*, Quantum deformations of algebras and their representations, Israel Math. Conf. Proc. 7, (1993)
- [8] G. Ginot, G. Halbout, *A formality theorem for Poisson manifold*, Letters in Mathematical Physics, **66**, (2003), 37–64.
- [9] G. Halbout, X. Tang, *Noncommutative Poisson structures on orbifolds*, arXiv: math.QA/0606436.
- [10] M. Kontsevich, *Deformation quantization of Poisson manifold*, Letters in Mathematical Physics, **66**, (2003), 157–216.
- [11] I. Moerdijk, *Orbifolds as groupoids: an introduction*, Adem, A. (ed.) et al., Orbifolds in mathematics and physics (Madison, WI, 2001), Amer. Math. Soc., Contemp. Math., 310, 205–222 (2002).
- [12] F. Nadaud, *Generalised Deformations, Koszul Resolutions, Moyal Products*, Reviews in Mathematical Physics, Vol. 10, No. 5 (1998) 685–704.
- [13] N. Neulaier, M. J. Pflaum, H. B. Posthuma and X. Tang, *Homology of formal deformations of proper éale Lie groupoids*, J. reine angew. Math. **593**, 117–168 (2006)
- [14] Serre, J., *Linear representations of finite groups*, Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42. Springer-Verlag, New York-Heidelberg, 1977.
- [15] P. Xu, *Noncommutative Poisson algebras*, Am. J. Math. 116, 101–125 (1994).

INSTITUT DE MATHÉMATIQUES ET DE MODÉLISATION DE MONTPELLIER, UNIVERSITÉ DE MONTPELLIER 2, CC5149, PLACE EUGÈNE BATAILLON, F-34095 MONTPELLIER CEDEX 5, FRANCE

E-mail address: halbout@math.univ-montp2.fr

INSTITUT DE MATHÉMATIQUES ET DE MODÉLISATION DE MONTPELLIER, UNIVERSITÉ DE MONTPELLIER 2, CC5149, PLACE EUGÈNE BATAILLON, F-34095 MONTPELLIER CEDEX 5, FRANCE

E-mail address: oudom@math.univ-montp2.fr

DEPARTMENT OF MATHEMATICS, WASHINGTON UNIVERSITY, ST. LOUIS, MISSOURI, USA, 63130

E-mail address: xtang@math.wustl.edu